

(19) World Intellectual Property
Organization
International Bureau




(43) International Publication Date
14 October 2004 (14.10.2004)

PCT

(10) International Publication Number
WO 2004/087892 A1

- (51) International Patent Classification⁷: C12N 1/20 (74) Agent: LEE, Won-Hee; 8th Fl. Sung-ji Heights II, 642-16
Yoksam-dong, Kangnam-ku, Seoul 135-080 (KR).
- (21) International Application Number: PCT/KR2004/000752 (81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG,
MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH,
PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (22) International Filing Date: 31 March 2004 (31.03.2004)
- (25) Filing Language: Korean
- (26) Publication Language: English
- (30) Priority Data:
10-2003-0020022 31 March 2003 (31.03.2003) KR
10-2003-0020023 31 March 2003 (31.03.2003) KR
- (71) Applicant (for all designated States except US): ALGEN-
TECH [KR/KR]; 599-1 Daeyeon-dong, Nam-gu, Busan
608-020 (KR).
- (71) Applicants and
(72) Inventors: -KIM, Young Tae [KR/KR]; #107-1002,
Kyungnamsunkyung Apt, Jwa-dong, Haeundae-gu, Bu-
san 612-030 (KR). LEE, Jae Hyung [KR/KR]; 167-2
Sambang-dong, Gimhae-si, Gyeongsangnam-do 621-190
(KR).
- (84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), Euro-
pean (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR,
GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
ML, MR, NE, SN, TD, TG).
- Published:
— with international search report
- For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: GENE INVOLVED IN THE BIOSYNTHESIS OF CAROTENOID AND MARINE MICROORGANISM, PARACOC-
CUS HAEUNDAENSIS, PRODUCING THE CAROTENOID



(57) Abstract: The present invention relates to a gene involved in the biosynthesis of carotenoid and a marine microorganism pro-
ducing the carotenoid, more particularly to a gene represented by Seq ID. No. 5, 7, 9, 11, 13 and 15 which encoding the protein
needed to biosynthesis of carotenoid and a marine microorganism, Paracoccus haeundaensis, producing the carotenoid. Since the
gene and the microorganism can effectively be used for, the massive-production of carotenoid.

WO 2004/087892 A1

11/PrA

GENE INVOLVED IN THE BIOSYNTHESIS OF
CAROTENOID AND MARINE MICROORGANISM,
PARACOCCLUS HAEUNDAENSIS, PRODUCING THE
CAROTENOID

5

FIELD OF THE INVENTION

The present invention relates to genes involved in the biosynthesis of carotenoid and a marine microorganism producing the carotenoid.

10

BACKGROUND


Carotenoid, a C40 isoprenoid compound having an anti-oxidant activity, means a group of pigment that is widely distributed in the nature. More than 600 kinds of carotenoids have been known so far, and they are all in different forms. The color of carotenoid varies from its molecular structure; that is whether it is yellow, red, scarlet or orange is decided upon the molecular structure of carotenoid. The examples of carotenoids are β -carotene (an orange pigment included in a carrot), lycopene (a red pigment included in a tomato), fucoxanthin (a yellowish brown or a brown pigment included in marine

20

plants), etc. As a precursor of vitamin A in human, carotenoid has activities of preventing oxidation, scavenging harmful oxygen, inhibiting the proliferation of cancer cells, and preventing the development of a cancer. It suggests that it has preventive effect on cardiovascular diseases, cancers and other adult diseases. It has been disclosed recently that carotenoid enhances immunity as being exposed on UV, so that it reduces skin damages by UV or inhibits the production of melanin. Since then, carotenoid came into a spotlight as a cosmetic material in Europe and in the U.S.A. Carotenoid is now in use as a health food ingredient (nutritional supplement), a pharmaceutical composition and a food-coloring agent, or as a pigment for animal feeds.

Among many carotenoids, astaxanthine (3, 3'-dihydroxy- β , β -carotene-4, 4'-dione) having the structure of below <Chemical Formula 1> is a scarlet or light orange color pigment produced in nature.

<Chemical Formula 1>



Astaxanthine is mostly included in tissues of marine animals such as shrimps, red seabreams, salmon and lobsters, etc (Fujita et al., *Nippon Suisan Gakkaishi.*, 49: 1855-1869, 1983; Johnson, E. A., *Crit. Rev. Biotechnol.*, 11: 297-326, 1991; Nelis et al., *J. Appl. Bacteriol.*, 70: 181-191, 1991). Astaxanthine not only inhibits the reactions of active oxygen to destroy DNA, proteins and lipids in cells during aerobic metabolism, to cause aging in cells and tissues, and to induce a cancer but also suppresses the generation of hydroxy or peroxy radicals (hydroxy or peroxy radicals (Palozza et al., *Arch. Biochem. Biophys.*, 297: 291-295, 1992; Shimidzu et al., *Fish Sci.*, 62: 134-137, 1996). In addition, astaxanthine has been known to have immune modulatory activity and cardioprotective effect (Jyonuchi et al., *Nutr. Cancer.*, 19: 269-280, 1993). In particular, an antioxidant activity of

astaxanthine is 10 times as high as that of other carotenoids and 100 times as high as that of α -tocopherol. However, toxicity of astaxanthine has not been reported as of today. Astaxanthine has
5 been widely used for the treatment and the prevention of various diseases including neurodegenerative diseases, cancers, immune disorders, cardiovascular diseases, etc, and studies are still going on further (Beal, H. F.,
10 *The Neuroscientist*, 3: 21-27, 1991; Chew et al., *Anticancer Res.*, 19: 1849-1853, 1999; Murillo E., *Arch. Latinoam. Nutr.*, 42: 409-413, 1992). Astaxanthine is also being industrially used as a coloring agent and has been registered as a food
15 additive in the name of 'Phaffia color' in Korea. The consumption of astaxanthine increases over 15% every year, suggesting the importance of astaxanthine.

A method for chemical synthesis of
20 astaxanthine has been recently developed by a company (Hoffman-LaRoche, Switzerland). However, a synthesized astaxanthine showed lower *in vivo* absorption and weaker stability as a food additive than a natural astaxanthine, so that the use of
25 the synthesized astaxanthine was allowed just in

some of European countries. Thus, a way to synthesize a natural astaxanthine is in strong demand and especially a way to produce astaxanthine using a microorganism producing
5 astaxanthine becomes the focus of industrial interest. *Phaffia rhodoxyma* (Miller et al., *Int. J. Syst. Bacteriol.*, 48: 529-536, 1976), a kind of yeast, *Haematococcus pluvialis* (Bubrick, *Bioresour Technol.*, 38: 237-239, 1991), a kind of
10 *Chlorophyta*, Gram-positive *Brevibacterium* 103 (Lizuka & Nishimura, *J. Gen. Appl. Microbiol.*, 15: 127-134, 1969), Gram-negative *Agrobacterium aurantiacum* (Yokoyama et al., *Biosci. Biotechnol. Biochem.*, 58: 1842-1844, 1994), *Paracoccus*
15 *marcusii* (Harker et al., *Int. J. Syst. Bacteriol.*, 48: 543-548, 1998) and *Paracoccus carotinifaciens* (Tsubokura et al., *Int. J. Syst. Bacteriol.*, 49: 277-282, 1999) are the examples of the microorganisms producing astaxanthine.

20

Studies have been focused on a gene coding an enzyme involved in biosynthesis of carotenoid for the past 6 years. As a result, a number of genes involved in biosynthesis of carotenoid were cloned
25 from various microorganisms and functions of them

were also examined (Armstrong, G. A., J. Bacteriol., 176: 4795-4802, 1994; Sandmann, G., Eur. J. Biochem., 223: 7-24, 1994; Wieland, B., J. Bacteriol., 176: 7719-7726, 1994). The pathway of
5 carotenoid biosynthesis is derived from FPP (farnesyl pyrophosphate), which is an intermediate product of general isoprenoid synthesis pathway. As seen in FIG. 8, FPP and IPP (isopentenyl pyrophosphate) turn into GGPP (geranylgeranyl
10 pyrophosphate) by geranylgeranyl pyrophosphate synthase encoded by *crtE*. Then, GGPP turns into β -carotene by the reactions of phytoene synthase encoded by *crtB*, phytoene desaturase encoded by *crtI* and lycopene cyclase encoded by *crtY*. β -
15 carotene changes into astaxanthine finally by the reactions of β -carotene ketolase encoded by *crtW* and β -carotene hydroxylase encoded by *crtZ*.

Nucleotide sequences, an organization and characteristics of *crt* gene (carotenogenic gene)
20 involved in biosynthesis of carotenoid have been investigated in *Rhodobacter capsulatus* (Armstrong et al., Mol. Gen. Genet., 216: 254-268, 1989), *Erwinia herbicola* (Sandimann et al., FEMS Microbiol. Lett., 71: 77-82, 1990; Hundle et al.,
25 Photochem. Photobiol., 54: 89-93, 1991) and

Ervinia uredovora (Misawa et al., *J. Bacteriol.*, 172: 6704-6712, 1990). Besides, *crt* gene involved in biosynthesis of carotenoid, which is composed of *crtB*, *crtI*, *crtY*, *crtW* and *crtZ*, has been
5 isolated from *Agrobacterium aurantiacum*, a marine microorganism (Norihiro et al., *J. Bacteriol.*, 177(22): 6575-6584, 1995). Another report has been made on three genes (*crtB*, *crtI* and *crtY*) coding enzymes catalyzing reactions from GGPP to
10 β -carotene and *Phaffia rhodozyma* where those three genes are inserted (WO 97/23633).

Thus, the present inventors isolated and identified a novel *Paracoccus* genus microorganism
15 producing astaxanthine and studied further to separate a gene involved in biosynthesis of carotenoid from the microorganism. As a result, the present inventors successfully cloned *crtE*, *crtB*, *crtI*, *crtY*, *crtW* and *crtZ* genes and *crt* gene
20 containing all of the above genes as well, and then the inventors examined nucleotide sequences of them, too. The present inventors completed this invention by confirming that carotenoid could be produced by using the *crt* gene in
25 microorganisms that were not able to produce

carotenoid.

SUMMARY OF THE INVENTION

It is an object of this invention to provide
5 a *Paracoccus haeundaensis* producing astaxanthine.

It is also an object of this invention to
provide a protein needed in biosynthesis of
carotenoid and a gene having nucleotide sequences
selected from a group consisting of nucleotide
10 sequences represented each by SEQ. ID. No 5, No 7,
No 9, No 11, No 13 and No 15.

It is a further object of this invention to
provide a gene involved in biosynthesis of
carotenoid containing the above gene. In
15 particular, the present invention provides a *crt*
gene having nucleotide sequences represented by
SEQ. ID. No 4.

It is another object of this invention to
provide a recombinant vector containing the gene
20 involved in biosynthesis of carotenoid above.

It is another object of this invention to
provide a method for producing carotenoid using
the above gene involved in biosynthesis of
carotenoid.

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

In order to achieve the above object, the present invention provides a *Paracoccus*
5 *haeundaensis* producing astaxanthine.

The present invention also provides a protein needed in biosynthesis of carotenoid and a gene having nucleotide sequences selected from a group consisting of nucleotide sequences each
10 represented by SEQ. ID. No 5, No 7, No 9, No 11, No 13 and No 15.

The present invention further provides a gene involved in biosynthesis of carotenoid containing the above gene. In particular, the present
15 invention provides *crt* gene having nucleotide sequences represented by SEQ. ID. No 4.

The present invention also provides a recombinant vector containing the gene involved in biosynthesis of carotenoid above.

20 The present invention further provides a method for producing carotenoid using the above gene involved in biosynthesis of carotenoid.

Hereinafter, the present invention is

described in detail.

The present invention provides a *Paracoccus haeundaensis* producing astaxanthine.

A microorganism was isolated from seawater sample taken from Haeundae shore in Busan, Korea as a strain having orange or red color. After investigating characteristics of the strain, it was confirmed that the strain was a rod type Gram-negative bacterium having non-motility and did not form spores (see FIG. 1). The cell size was 0.3-0.7 μm in diameter and 0.8-2.5 μm in length. The colony had orange (scarlet) color. The optimum growth temperature of the strain was 25°C, and the strain was never growing under 10°C or over 40°C (see FIG. 2). The optimum NaCl concentration for the growth was 1-6% (w/w) and the strain was not growing at all with over 8% (see FIG. 4). The optimum growth pH of the strain was 8 (see FIG. 3).

The strain of the present invention used only D-arabinose and galactose as a carbon source and an energy source for the growth. Neither pentoses, hexoses, sugar alcohols, oligosaccharides nor other amino acids were used. Starch hydrolysis, cytochrome oxidase and catalase reactions were all positive. But, urease reaction was negative. The

strain could not produce indole from tryptophane, but took advantage of citric acid in deed, confirmed by citric acid test. Denitrification test was also performed. As a result, the strain
5 reduced nitrate to nitrite, but did not reduce nitrite to N₂ gas. Besides, the strain did not ferment glucose. The strain of the present invention was also confirmed to be aerobic. DNA G+C composition of the strain was 66.9 mol%. The
10 major non-hydroxyl fatty acid was unsaturated C18:1, and the major hydroxyl fatty acid was C10:0 (3-OH).

In general, a *Paracoccus* genus microorganism is an oxidase and catalase positive, gram-negative
15 bacterium and belongs to α -3-subclass of *Proteobacteria* phylogenetically. Other characteristics of the microorganism were also investigated (see Table 2). As a result, it was confirmed that the strain of the present invention
20 belonged to *Proteobacteria*.

In order to identify the strain of the present invention more concretely, the inventors examined sequence of 16S rDNA. As a result, the
25 nucleotide sequence of 16S rDNA of the strain of

the invention had a high homology with those of *Paracoccus marcusii* and *Paracoccus carotinifaciens*. Nevertheless, the strain of the present invention showed different characteristics, comparing to other *Paracoccus* genus microorganisms including the two above (see Table 4 and Table 5). Therefore, the present inventors confirmed that the strain of the invention was a novel *Paracoccus* genus microorganism and named it '*Paracoccus haeundaensis*'.

The present inventors investigated if carotenoid was generated in the *Paracoccus haeundaensis*, resulting in the confirmation that the strain of the present invention produced β -carotene and astaxanthine (see FIG. 7). The present inventors deposited the *Paracoccus haeundaensis* of the present invention at KCCM (Korean Culture Center of Microorganisms) on January 24, 2003 (Accession No: KCCM - 10460).

Paracoccus haeundaensis of the present invention can be effectively used for the production of carotenoid, especially astaxanthine. A method to produce carotenoid using a microorganism of the present invention includes

the steps of culturing *Paracoccus haeundaensis* in a proper medium and collecting astaxanthine from the culture solution. Particularly, the strain of the present invention was cultured in PPES-II
5 medium at 25°C for 4 days, which was the primary culture, then cells were collected from the culture solution. Organic solvent was added to the cells, which were cultured at 4°C for overnight, resulting in the elution of
10 astaxanthine. In addition to the above PPES-II medium, LB3 medium (LB medium complemented with 3% NaCl) can be used as a medium for the culture of the strain. It is also preferable to add galactose to a medium in order to induce
15 satisfactory growth of the strain and mass-produce astaxanthine. Methanol, acetone or ethyl ether can be used as an organic solvent used for the purification of astaxanthine from the cultured cells, and methanol is more preferably used. The
20 collection of astaxanthine from the cultured cells can be performed using HPLC (high performance liquid chromatography) or TLC (thin-layer chromatography) by the conventional method known to the people in this field.

25

The present invention also provides a protein required for biosynthesis of carotenoid and a gene having nucleotide sequences selected from a group consisting of sequences each represented by SEQ. ID. No 5, No 7, No 9, No 11, No 13 and No 15.

The gene above means a gene producing astaxanthine purified from *Paracoccus haeundaensis* (Accession No: KCCM-10460).

It is preferred for a gene coding a protein required for biosynthesis of carotenoid of the present invention to have a sequence selected from a group consisting of sequences each represented by SEQ. ID. No 5, No 7, No 9, No 11, No 13 and No 15. Each protein coded by each gene above has amino acid sequence represented by SEQ. ID. No 6, No 8, No 10, No 12, No 14 and No 16, respectively. 6 genes of the present invention and proteins coded by the same are shown in the below Table 1.

<Table 1>

Gene	Gene name	Protein	Amino acid sequence
SEQ. ID. No 5	<i>crtW</i>	β -carotene ketolase	SEQ. ID. No 6
SEQ. ID. No 7	<i>crtZ</i>	β -carotene hydroxylase	SEQ. ID. No 8

SEQ. ID. No 9	<i>crtY</i>	Lycopene cyclase	SEQ. ID. No 10
SEQ. ID. No 11	<i>crtI</i>	Phytoene desaturase	SEQ. ID. No 12
SEQ. ID. No 13	<i>crtB</i>	Phytoene synthase	SEQ. ID. No 14
SEQ. ID. No 15	<i>crtE</i>	Geranylgeranyl pyrophosphate synthase	SEQ. ID. No 16

Genes provided by the present invention can be effectively used for the production of carotenoid by being inserted in various host cells.

5 Those genes can be used either singly or together (more than 2, at least). For example, a gene coding lycopene cyclase and represented by SEQ. ID. No 9 can be used for the production of β -carotene by being inserted in a microorganism containing

10 *crtE*, *crtB* and *crtI* only. And, genes represented by SEQ. ID. No 5 and No 7, coding β -carotene ketolase and β -carotene hydroxylase respectively, can be used for the production of astaxanthine by being inserted in a microorganism producing β -

15 carotene (ex: *Phaffia rhodozyma* ATCC96815).

The present invention further provides a carotenoid synthesis gene containing all the above genes.

20 A carotenoid synthesis gene is preferred to

have nucleotide sequences represented by SEQ. ID. No 4. The carotenoid synthesis gene (referred as 'crt gene' hereinafter) of the present invention includes all the carotenoid synthesis genes involved in astaxanthine production process. An organization of crt gene of the present invention is presented in FIG. 15. As shown in FIG. 15, the size of crt gene of the present invention is 6,223 bp and includes *crtW*, *crtZ*, *crtY*, *crtI*, and *crtB* in that order in 5'→ 3' direction. Each of *KpnI*, *SmaI*, *XmaI*, *ClaI*, *HindIII* and *BamHI* recognition sequence is located therein. Stop codon of each *crtW*, *crtZ*, *crtY*, *crtI* and *crtB* is overlapped to start codon of the next gene. In particular, *crtE* gene is found as a complementary strand.

The present invention also provides a recombinant vector containing the above carotenoid biosynthesis gene.

A recombinant vector of the present invention was constructed by inserting crt gene into a basic vector. Any basic vector that was generally used for gene cloning or expression could be used for the present invention without limitation. And a choice of a vector depended on a host cell. For

example, if *E. coli* is used as a host cell, an *E. coli* specific vector having replication origin of the *E. coli* is preferred. Likewise, if yeast is used as a host cell, a yeast specific vector
5 having replication origin of yeast is preferred. A shuttle vector that has both replication origin of *E. coli* and replication origin of yeast at the same time is also available. In the preferred embodiment of the present invention, the present
10 inventors constructed a recombinant vector containing *crt* gene by using pCR-XL-TOPO vector, which was named 'pCR-XL-TOPO crtfull'.

The present invention also provides a strain
15 prepared by transformed host cells with a recombinant vector containing the above carotenoid biosynthesis gene.

E. coli or yeast can be used as host cells of the present invention, and in particular, *E. coli*
20 is preferably selected from a group consisting of XLI-Blue, TOPO, BL21(DE3) codon plus, DH1 and DH5a, but the choice is not always limited thereto. In a preferred embodiment of the present invention, a strain was prepared by transformed
25 BL21(DE3) codon plus, a kind of *E. coli*, with a

recombinant vector 'pCR-XL-TOPO crtfull' which contained crt gene represented by SEQ. ID. No 4.

5 The present invention further provides a method for the production of carotenoid using the carotenoid biosynthesis gene.

The carotenoid producing method of the present invention is comprised of the following steps:

10 1) Cloning crt gene represented by SEQ. ID. No 4;

2) Constructing a recombinant vector in which the gene of the above step 1) was inserted;

15 3) Transforming a host cell with the recombinant vector of the step 2); and

4) Recovering carotenoids from the culture solution in which a strain transformed with the above recombinant vector was being cultured.

20 *E. coli* can be used as a host cell. At this time, any *E. coli* strain generally used for the transformation can be used without limitation, but it is preferred to select *E. coli* from a group consisting of XLI-Blue, TOPO, BL21(DE3) codon plus, DH1 and DH5a. In a preferred embodiment of the present invention, BL21(DE3) codon plus was

25

selected. The choice of a host cell for the invention is not limited to *E. coli*, and yeast is also available.

Particularly, a strain constructed in the
5 above step 1) ~ step 3) was cultured in a growth medium (primary culture). Cells were recovered from the culture solution. Organic solvent was added to the cells, which was further cultured at 4°C for overnight (secondary culture). At that
10 time, it was possible to add IPTG (isopropyl-beta-D-thiogalactopyranside), an inducer inducing the production of carotenoid, into the culture solution. Carotenoid substrates such as FPP (farnesyl pyrophosphate), GGPP (geranylgeranyl
15 diphosphate) or GPP (geranylpyrophosphate) could also be added. Methanol, acetone or ethyl ether could be used as an organic solvent for the extraction of carotenoid from the culture cells, and methanol was more preferred. Carotenoid was
20 recovered from the culture cells by using HPLC (high performance liquid chromatography) or TLC (thin-layer chromatography) by following the conventional method commonly known for the people in this field.

25

In a preferred embodiment of the present invention, the inventors measured the amount of astaxanthine produced from a transgenic strain containing *crt* gene. As a result, the produced
5 astaxanthine was 110 $\mu\text{g/g}$ (dry weight), which was far more than that produced by an astaxanthine producing strain '*Paracoccus haeundaensis*' (25 $\mu\text{g/g}$ (dry weight)). Therefore, the method for producing astaxanthine of the present invention
10 makes possible even for a strain which cannot produce astaxanthine itself to mass-produce astaxanthine by using carotenoid biosynthesis gene, so that it facilitates the production of medical supplies and edible pigments as a food additive
15 containing astaxanthine.

In an example of the present invention, a genomic DNA library of *Paracoccus haeundaensis* was constructed in order to clone a gene coding a
20 protein required for carotenoid biosynthesis. The construction of a genomic DNA library was performed by the conventional method commonly known to the people in this field. In particular, a genomic DNA library was constructed by using a
25 cosmid vector in this invention.

In another example of the present invention, 'color complementation' was used for cloning genes, involved in carotenoid biosynthesis, from a genomic DNA library. A microorganism not
5 producing carotenoid (ex: *E. coli*) is given power to generate carotenoid by being transformed with a carotenoid biosynthesis gene that was cloned from a carotenoid producing microorganism (ex: *Paracoccus haeundaensis* of the present invention).
10 For example, *E. coli* could produce β -carotene and cells turned into yellow after being transformed with *crtE*, *crtB*, *crtI* and *crtY*. And *E. coli* transformed with *crtE*, *crtB*, *crtI*, *crtY*, *crtW* and *crtZ* could produce astaxanthine and cells turned
15 into orange. The present inventors cultured a genomic DNA library of *Paracoccus haeundaensis* in a medium supplemented with FPP (farnesyl pyrophosphate), a common substrate of carotenoid. As a result, 13 colonies having orange color were
20 selected. Then, a cosmid vector was isolated from each colony. Nucleotide sequence of DNA insert included in the vector was identified. As a result, it was confirmed that the size of the smallest DNA insert was 6,223 bp. The nucleotide
25 sequence of the DNA insert was represented by SEQ.

ID. No 4.

In another example of the present invention, nucleotide sequences of 6,223 bp long DNA insert assumed to contain a gene producing carotenoid was investigated, resulting in the obtainment of 6 ORFs. Those were analyzed nucleotided on NCBI GenBank. As a result, amino acid sequences translated from each ORF had a high homology with amino acid sequences of 6 enzymes involved in the reaction inducing astaxanthine production from FPP (see FIG. 9 - FIG. 14). From the result, the present inventors confirmed that the DNA insert isolated in this invention had *crt* gene coding a protein necessary for carotenoid biosynthesis (see FIG. 15).

In another example of the present invention, the inventors investigated if *E. coli* not producing carotenoid could produce it by the insertion of *crt* gene coding a relevant protein. First, recombinant vector 'pCR-XL-TOPO-crtfull', in which *crt* gene was inserted, was constructed (see FIG. 16). Then, the prepared recombinant vector was inserted in *E. coli*. Lastly, *E. coli* transformant having orange color was selected. In order to confirm if the transformant could produce

astaxanthine, the transformant was cultured and then cells were collected. Methanol was added to the obtained cells, which were cultured at 4°C for overnight. Then, supernatant was obtained and optical density was measured at 190-900 nm. As a result, as shown in FIG. 17, original peaks of β -carotene and astaxanthine were confirmed. For more accurate analysis, HPLC assay was performed with some of the supernatant. At that time, β -carotene and astaxanthine purchased from Sigma were used as standard substances. As a result, it was confirmed that a transformant, in which *crt* gene isolated by the present inventors was inserted, could produce β -carotene and astaxanthine. The amount of astaxanthine produced from the transformant of the present invention was 110 $\mu\text{g/g}$ (dry weight). The result suggested that *E. coli* in which *crt* gene was inserted could produce astaxanthine far more than *Paracoccus haeundaensis*, an astaxanthine producing strain, could do (25 $\mu\text{g/g}$ (dry weight)).

BRIEF DESCRIPTION OF THE DRAWINGS

The application of the preferred embodiments

of the present invention is best understood with reference to the accompanying drawings, wherein:

FIG. 1 is a set of photographs showing the
5 result of observation with a transmission electron microscope on the strain of the present invention in exponential stage, and each bar is 200 nm in length.

10 FIG. 2 is a graph showing the growth curve of the strain of the present invention depending on temperature.

FIG. 3 is a graph showing the growth curve of
15 the strain of the present invention depending on pH.

FIG. 4 is a graph showing the growth curve of
the strain of the present invention depending on
20 culture time.

FIG. 5 is a graph showing the growth curve of
the strain of the present invention depending on
NaCl concentration.

25

FIG. 6 is a schematic diagram showing the phylogenetic dendrogram of the *Paracoccus* genus microorganisms of the present invention, based on 16S rDNA sequence data, and a bar represents maximum-parsimony distance (1 nucleotide substitutions per 100 nucleotides).

FIG. 7 is a set of graphs showing the results of scanning on the ranges of optical density changes at 190 ~ 890 nm (A) and at 400 ~ 550 nm (B) of a methanol extract extracted from the culture solution of the strain of the present invention. The peak at 450 nm is the peculiar peak of β -carotene, and the peak at 470 nm is the original peak of astaxanthine.

FIG. 8 is a schematic diagram showing the pathway of carotenoid biosynthesis.

FIG. 9 is a schematic diagram showing the result of comparison between an amino acid sequence of the first ORF (open reading frame) included in a DNA insert isolated from *Paracoccus haeundaensis* and amino acid sequences of β' -carotene ketolase (crtW) isolated from an

Alcaligenes genus microorganism and a
Bradyrhizobium genus microorganism.

P. haeundaensis: Paracoccus haeundaensis

5 *Alcaligenes*_sp: *Alcaligenes* genus
microorganism

*Bradyrhizobium*_sp: *Bradyrhizobium* genus
microorganism

Consensus: corresponding amino acid sequence

10 FIG. 10 is a schematic diagram showing the
result of comparison between amino acid sequences
of the second ORF included in a DNA insert
isolated from *Paracoccus haeundaensis* and amino
acid sequences of β -carotene hydroxylase (*crtZ*)
15 isolated from an *Alcaligenes* genus microorganism.

P. haeundaensis: Paracoccus haeundaensis

*Alcaligenes*_sp: *Alcaligenes* genus
microorganism

Consensus: corresponding amino acid sequence

20

FIG. 11 is a schematic diagram showing the
result of comparison between amino acid sequences
of the third ORF included in a DNA insert isolated
from *Paracoccus haeundaensis* and amino acid
25 sequences of lycopene cyclase (*crtY*) isolated from

an *Flavobacterium* genus microorganism.

P. haeundaensis: *Paracoccus haeundaensis*

Flavobacterium_sp: *Flavobacterium* genus
microorganism

5 Consensus: corresponding amino acid sequence

FIG. 12 is a schematic diagram showing the
result of comparison between amino acid sequences
of the fourth ORF included in a DNA insert
10 isolated from *Paracoccus haeundaensis* and amino
acid sequences of phytoene desaturase (*crtI*)
isolated from an *Flavobacterium* genus
microorganism.

P. haeundaensis: *Paracoccus haeundaensis*

15 Flavobacterium_sp: *Flavobacterium* genus
microorganism

Consensus: corresponding amino acid sequence

FIG. 13 is a schematic diagram showing the
20 result of comparison between amino acid sequences
of the fifth ORF included in a DNA insert isolated
from *Paracoccus haeundaensis* and amino acid
sequences of phytoene synthase (*crtB*) isolated
from an *Flavobacterium* genus microorganism.

25 P. haeundaensis: *Paracoccus haeundaensis*

Flavobacterium_sp: *Flavobacterium* genus
microorganism

Consensus: corresponding amino acid sequence

5 FIG. 14 is a schematic diagram showing the
result of comparison between amino acid sequences
of the sixth ORF included in a DNA insert isolated
from *Paracoccus haeundaensis* and amino acid
sequences of geranylgeranyl pyrophosphate synthase
10 (*crtE*) isolated from an *Flavobacterium* genus
microorganism.

P. haeundaensis: Paracoccus haeundaensis

Flavobacterium_sp: *Flavobacterium* genus
microorganism

15

FIG. 15 is a schematic diagram showing the
organization of *crt* gene isolated from *Paracoccus*
haeundaensis.

20

FIG. 16 is a schematic diagram showing the
cleavage map of pCR-XL-TOPO-*crt*full vector of the
present invention.

FIG. 17 is a set of graphs showing the
25 results of scanning on the ranges of optical

density changes at 190 ~ 890 nm (A) and at 350 ~ 550 nm (B) of a methanol extract extracted from the culture cells of the transformant of the present invention in which crt gene was inserted.

- 5 The peak at 450 nm is the peculiar peak of β -carotene, and the peak at 470 nm is the original peak of astaxanthine.

EXAMPLES

- 10 Practical and presently preferred embodiments of the present invention are illustrative as shown in the following Examples.

However, it will be appreciated that those skilled in the art, on consideration of this
15 disclosure, may make modifications and improvements within the spirit and scope of the present invention.

- Example 1: Sample collection, microorganism
20 separation, cultivation and maintenance of the
same

Seawater sample taken from Haeundae shore in

Busan, Korea, was diluted by 1/1000, which was then smeared on a nutrient agar medium (Difco) and cultured at 25°C for 3 days. Among microorganisms cultured thereby, those having orange or red color were isolated. In order to isolate a microorganism producing carotenoid among strains isolated, all the candidates were cultured in PPES-II medium (tripton 1 g/l, bacto-soyton 1 g/l, ferric citrate 0.01 g/l, polypeptone 2 g/l and sodium chloride 3 g/l). Cells were collected from the culture solution. Carotenoid was extracted by using methanol to identify a final strain producing carotenoid. The isolated microorganism was named 'BC74171^T strain'.

15

Example 2: Investigation of phenotypic characteristics

In order to investigate morphological and physiological characteristics of the BC74171^T strain of the present invention, experiments were performed as follows.

20

<2-1> Morphological characteristics

BC74171^T strain of the present invention was cultured in PPES-II medium at 25°C for 3 days. The cultured cells were suspended in 0.1 M phosphate buffer (pH 7.2). The cells were fixed
5 by 2% glutaraldehyde, washed with 0.05 M cacodylate buffer solution, and fixed again by 1% osmium tetroxide. The fixed cells were dehydrated in ethanol that was later replaced with propylene. Those cells were put into EPON resin and sections
10 were prepared using an ultramicrotome. The prepared sections were observed under a JEM 1200EX-II transmission electron microscope (TEM).

As a result, as shown in FIG. 1, BC74171^T strain of the present invention had a shape of rod
15 and was 0.3-0.7 μm in diameter and 0.8-2.5 μm in length. The strain did not form spores. The mobility of the strain was also observed under an optical microscope by hanging-drop technique (Skerman, V. B. D., *A Guide to the Identification*
20 *of the Genera of Bacteria*, 2nd edn. Baltimore, 1967). As a result, BC74171^T strain of the present invention was non-motile. In addition, BC74171^T strain was Gram-negative, and its colonies were flat and had a light orange color.

<2-2> Physiological characteristics

1. Range of temperature

BC74171^T of the present invention was cultured in a nutrition agar medium (Difco) at different temperatures (4, 10, 20, 25, 28, 30, 37, 40 and 50°C) for 10 days to investigate the range of growth temperature. As a result, BC74171^T strain was growing at temperature ranging 20-37°C, and the optimum growth temperature was 25°C (Table 2 and FIG. 2).

<Table 2>

Characteristics of *Paracoccus haeundaensis*

Morphological characteristics		Utilization	
Cell diameter (μm)	0.3-0.7	D-Glucose	—
Cell length (μm)	0.8-2.5	Maltose	—
Mobility	*—	D-Galactose	*+
Spore formation	—	Sucrose	—
Optimum growth temperature (°C)	25	Mannitol	—
Optimum growth pH	8	Cellobiose	—
NaCl resistance (%)	7	Trehalose	—

Product		Xylose	—
Indole	—	Dulcitol	—
MR (Methyl red test)	—	Salicin	—
VP (Voges-Proskauer test)	—	Adonitol	—
Hydrogen sulfide	—	Inositol	—
Citrate	+	Arabinose	+
Enzyme activity		Raffinose	—
Catalase	+	Rhamnose	—
Urease	—	D-Fructose	—
Oxidase	+	D-Mannose	—
Starch hydrolysis	+	Dimethylformamide	—
Denitrification		Glycerol	—
Nitrate → Nitrite	+	L-Glutamic acid	—
Nitrite → N ₂ gas	—	Sorbitol	—
Color	+(light orange)	Lactose	—
Chemotaxonomic characteristics		L-Asparagine	—
G + C content (mol%)	66.9	Acetone	—
Non-hydroxyl acids	C _{18:1}	Major carotenoid	Astaxanthine
3-Hydroxyl fatty acids	3-OH C _{10:0}		

*—: Positive response,

*+: Negative response

2. Range of pH

BC74171^T of the present invention was cultured in PPES-II media each having different pH (3.0, 4.0, 5.0, 6.0, 7.0, 8.0, 9.0, 10.0, 10.5 and 11.0) for 10 days to investigate growth pH range.

5 As a result, the optimum growth pH was 8. The growth of the strain was inhibited or retarded under pH 6 and over pH 10.5 (Table 2 and FIG. 3).

3. Growth

10 BC74171^T strain of the present invention was shaking-cultured in a PPES-II medium at 25°C for 10 days to investigate its growth by measuring turbidity. As a result, the growth of the strain was rapidly increased from 30 hours after the
15 culture and slowly decreased from 50 hours after the culture (FIG. 4).

4. Resistance against NaCl

BC74171^T of the present invention was
20 cultured in a trypticase soy broth medium containing 1-10%(w/v) NaCl for 10 days at various temperatures (4, 10, 20, 25, 28, 30, 37, 40 and 50°C). As a result, the optimum NaCl concentration for growth was 1-6%. The growth was
25 retarded with 7% NaCl and was stopped over 8%

(Table 2 and FIG. 5).

5. Utilizing ability of carbon source

In order to examine a capacity to use a carbon source, a micro-plate including 24 substrates listed in the below Table 3 was used. The final concentration of each substrate was adjusted to 1%, which was loaded in a purple broth base (Difro), after being filtered or sterilized by moist heat. The medium was inoculated with the strain of the present invention, then was cultured at 25°C for 10 days. Yellow meant a positive reaction and purple, which was an original color of the medium, meant a negative reaction.

<Table 3>

Substrates used in the carbon source utilizing ability test and the utilizing ability

	Substrate	Utilizin g ability		Substrate	Utilizin g ability
1	D-Glucose	*-	13	Arabinose	+
2	Maltose	-	14	Raffinose	-
3	D-Galactose	*+	15	Rhamnose	-
4	Sucrose	-	16	D- Fructose	-
5	Mannitol	-	17	D-Mannose	-

6	Cellobiose	-	18	Dimethylformamide	-
7	Trehalose	-	19	Glycerol	-
8	Xylose	-	20	L-Glutamic acid	-
9	Dulcitol	-	21	Sorbitol	-
10	Salicin	-	22	Lactose	-
11	Adonitol	-	23	L-Asparagine	-
12	Inositol	-	24	Acetone	-

*-: Not used (positive reaction),

+: Used (negative reaction)

As a result, as shown in Table 3, BC74171^T strain of the present invention used only D-arabinose and galactose as a carbon source, and no other substrates were used.

6. Activity of starch hydrolysis

In order to investigate if the strain of the present invention could hydrolyze starch, a starch agar was inoculated with the strain, then, cultured for 10 days. An activity of starch hydrolysis was measured by the method of Cowan, S. T. & Steel, K. J. (Cowan, S. T. & Steel, K. J., *Manual for the Identification of Medical Bacteria*. London: Cambridge University Press, 1965). As a

result, the strain of the present invention was confirmed to have an activity of starch hydrolysis (Table 2).

5 7. Indole test (tryptophanase activity)

 In order to investigate if the strain of the present invention could produce indole by decomposing tryptophane, the present inventors took advantage of the method of Cappuccino et al. (Cappuccino J. G. and Sherman, N. In *Microbiology: a laboratory manual* (6th) 2001), resulting in the confirmation that indole was not produced from tryptophane (Table 2).

15 8. Acid production (fermentation) from
carbohydrates

 Acid production was investigated by the method of Hughm et al. (Hughm et al., *J. Bacteriol.*, 66: 24-26, 1953). Methyl Red test and
20 Voges-Proskauer test were performed and the results were all negative (Table 2).

9. Citric acid test

 In order to investigate if the strain of the present invention could be growing by using citric
25 present invention could be growing by using citric

acid as a carbon source, the method of Cappuccino et al. (Cappuccino J. G. and Sherman, N. In *Microbiology: a laboratory manual (6th)* 2001) was performed and the result was positive (Table 2).

5

10. Catalase activity

The catalase activity was investigated by observing bubbles in 3% hydrogen peroxide solution by following the method of Harker et al. (Harker et al., *J. Clin. Microbiol.*, 2: 463-464, 1975). As a result, the catalase activity of the strain of the present invention was confirmed to be positive (Table 2).

15 11. Oxidase activity

The oxidase activity was investigated by using 1% p-aminodimethylaniline oxalate as a substrate by following the method of Cappuccino et al. (Cappuccino J. G. and Sherman, N. In *Microbiology: a laboratory manual (6th)* 2001). As a result, the cytochrome oxidase activity of the strain of the present invention was confirmed to be positive (Table 2).

25 12. Urease activity

The urease activity was measured by the method of Lanyi (Lanyi, B. *Methods Microbiol.*, 19: 1-67, 1987). As a result, the urease activity of the strain of the present invention was confirmed to be negative (Table 2).

13. Denitrification

In order to observe denitrification, gas production and growth were investigated through stab culture using a nutritive medium containing 0.1% (w/v) agar, by following the method of Cappuccino et al. (Cappuccino J. G. and Sherman, N. In *Microbiology: a laboratory manual (6th)* 2001). As a result, BC74171^T strain of the present invention reduced nitrate to nitrite. But, the strain could not reduce nitrite to N₂ gas (Table 2).

14. Hydrogen sulfide production test

In order to investigate if the strain of the present invention could produce hydrogen sulfide from amino acid containing sulfur like cysteine or a substrate like inorganic sulfur compounds, hydrogen sulfide production test was performed using TSI (Triple Sugar Iron Agar) medium

(Cappuccino J. G. and Sherman, N. In *Microbiology: a laboratory manual* (6th) 2001). The result was negative (Table 2).

5 Example 3: Fatty acid analysis

BC74171^T strain of the present invention was cultured on trypticase soy agar medium (pH 8.0) supplemented with 2% NaCl at 25°C for 2 days. Then, harvesting, saponification, methylation and
10 extraction of FAMES (fatty acid methyl esters) were performed with the cells cultured by the method of Sasser (Sasser, M. In *Methods in Phytobacteriology*, 199-204, 1990). GC-MS was performed following the method of Lipske et al.
15 (Lipski et al., *Syst. Appl. Microbiol.*, 20: 448-457, 1997), leading to the investigation of FAMES. As a result, according to the confirmed fatty acid profile ($C_{18:1}$, 84.32 %; $C_{18:0}$, 7.79 %; $C_{10:0}$ (3-OH), 2.06 %; $C_{12:1cis5}$ 2.0 %; $C_{14:0}$ (3-OH), 1.47 %; $C_{17:0}$,
20 0.80 %; $C_{16:0}$, 0.78 %; and unknown peak, 0.78 %), the strain of the present invention was confirmed to belong to α -subclass of *Proteobacteria*. And a major hydroxyl fatty acid was $C_{10:0}$ (3-OH) (Table 2).

Example 4: Base composition of DNA

The base composition of genomic DNA of BC74171^T strain of the present invention was identified by the method of Tamaoka et al. (Tamaoka, J. & Komagata, K. *FEMS Microbiol. Lett.*, 25: 125-128, 1984). Particularly, the genomic DNA was extracted from BC74171^T strain of the present invention by following the conventional method known to the people in this field. The extracted genomic DNA was hydrolyzed. The obtained nucleotides were analyzed by HPLC (reverse-phase HPLC). As a result, G+C content of DNA of BC74171^T strain was 66.9 mol% (Table 2).

15

Example 5: 16S rDNA sequencing and phylogenetic dendrogram analysis

Genomic DNA was extracted from BC74171^T strain of the present invention by the method of Rainey et al. (Rainey et al., *Syst. Appl. Microbiol.*, 15: 197-202, 1992). 16S rDNA was PCR amplified by using primers represented by SEQ. ID. No 1 and no 2. PCR was performed as follows.

20

pmol each of two primers, 10 ng of genomic DNA, 1 unit of Taq polymerase and 10x buffer solution (with $MgCl_2$) were mixed to make the reacting solution. The prepared reacting solution was pre-denatured at 94°C for 5 minutes, followed by 25 cycles of denaturation at 94°C for 1 minute, annealing at 56°C for 30 seconds, polymerization at 72°C for 90 seconds, and final extension at 72°C for 10 minutes. The PCR product was cloned into pGEM-T vector (Promega) and isolated DNA clones were reacted with ABI PRISM™ staining reagent (Perkin Elmer, USA). Nucleotide sequence was determined by using ABI 377 genetic analyzer (Perkin Elmer, USA).

As a result, 16S rDNA of BC74171^T strain of the present invention was 1451 bp long and had nucleotide sequences represented by SEQ. ID. No 3. The nucleotide sequence of 16S rDNA of BC74171^T strain was analyzed by using BLASTN and BLASTX of NCBI GenBank database. As a result, the nucleotide sequence of 16S rDNA of BC74171^T strain showed 99.8% and 99.6% homology each with those of *Paracoccus marcusii* and *Paracoccus carotinifaciens* which have been known to produce astaxanthine among various *Paracoccus* genus microorganisms.

Thus, BC74171^T strain of the present invention was confirmed to be a *Paracoccus* genus strain. And the characteristics of the strain of the present invention were compared with those of various *Paracoccus* genus microorganisms. The characteristics of other *Paracoccus* genus microorganism listed in the below Table 4 (2-15) were the results of investigations by Harker et al. (Harker et al., *Int. J. Syst. Bacteriol.*, 48: 543-548, 1998), Lipski et al. (Lipski et al., *Syst. Appl. Microbiol.*, 20: 448-457, 1997), Tsubokura et al. (Tsubokura et al., *Int. J. Syst. Bacteriol.*, 49: 277-282, 1999), Kelly et al. (Kelly et al., *The genus Paracoccus. In The Prokaryotes*, <http://www.prokaryotes.com>., Edited by M. Dworkin, S. Falkow, E. Rosenberg, K. H. Schleifer & E. Stackebrandt. New York: Springer., 2000) and Doronina et al. (Doronina et al., *Int. J. Syst. Evol. Bacteriol.*, 52: 679-6, 2002). As shown in Table 4, BC74171^T strain of the present invention did not share any characteristics with any other *Paracoccus* genus microorganism.

<Table 4>

Comparison of the characteristics of the strain of

the present invention and those of other
Paracoccus genus microorganisms

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
Mobility	-	-	-	-	-	+	-	-	-	-	-	-	-	-	+
Growth in 6% NaCl	+	-	NR	-	-	NR	-	-	-	W	NR	NR	-	NR	NR
Usefulness	Glucose	-	+	-	+	+	+	-	+	+	+	+	-	+	+
	Arabinose	+	+	-	-	-	-	-	+	+	+	-	-	-	-
	Glycerol	-	+	-	+	+	NR	+	-	+	+	+	-	-	+
	Sucrose	-	-	-	-	-	NR	+	-	-	+	+	+	-	+
	Mannitol	-	+	-	-	+	+	+	-	+	+	+	+	-	+
	Fructose	-	+	-	-	+	NR	+	-	+	+	+	+	-	+
Denitri- fication	-	-	+	-	-	-	+	+	+	-	-	+	+	+	+
Urease activity	-	+	NR	-	-	-	+	-	+	NR	+	-	NR	-	-
Yellow color	+	-	NR	-	-	+	-	-	-	+	-	-	-	-	-
G + C content (mol%)	66.9	64-66	NR	63	67	67	64-67	71	62.5	66	67	66	68.5-70	66-67.6	67-68

*1: The strain of the present invention

5 2: *P. alcaliphilus* (JCM 7364T)

- 3: *P. alkenifer* (DSM 11593T)
 4: *P. aminophilus* (JCM 7686T)
 5: *P. aminovorans* (JCM 7685T)
 6: *P. carotinifaciens* (IFO 16121T)
 5 7: *P. denitrificans* (ATCC 17741T)
 8: *P. kocurii* (JCM 7684T)
 9: *P. kondratievae* (VKM B-2222T)
 10: *P. marcusii* (DSM 11574T)
 11: *P. methylutens* (VKM B-2164T)
 10 12: *P. pantotrophus* (ATCC 35512T)
 13: *P. solventivorans* (DSM 6637T)
 14: *P. thiocyanatus* (IAM 12816T)
 15, *P. versutus* (ATCC 25364T)
 *+: Positive reaction -: Negative reaction
 15 W: Weak reaction) NR: Not reported

The characteristics of BC74171^T strain of the present invention were different in many ways even with *Paracoccus marcusii* and *Paracoccus carotinifaciens* showing the highest 16S rDNA sequence homology. More precisely, as shown in the below Table 5, *Paracoccus marcusii* (Harker et al., *Int. J. Syst. Bacteriol.*, 48: 543-548, 1998) was a coccus or a short rod type forming a short chain. On the contrary, BC74171^T strain of the

present invention was a rod type not forming a chain. *Paracoccus carotinifaciens* (Tsubokura et al., *Int. J. Syst. Bacteriol.*, 49: 277-282, 1999) was also a rod type not forming a chain but had flagella. However, BC74171^T strain of the present invention had not flagella. The strain of the present invention did not use any of glucose, mannitol, maltose or mannose, but the other two strains did use them. Besides, BC74171^T strain of the present invention decomposed starch. But, the two other strains did not.

<Table 5>

Comparison of the characteristics of the strain of the present invention and those of other *Paracoccus* genus microorganisms

Characteristics	BC74171 ^T	<i>Paracoccus marcusii</i>	<i>Paracoccus carotinifaciens</i>
Cell morphology	Rod type	Coccus or short rod type	Rod type
Mobility	- *	-	+ *
Uses of ul- ne- s s	Glucose	-	+
	Maltose	-	+
	Mannitol	-	+
	Arabinose	+	-

Citric acid	+	+	-
Mannose	-	NR*	+
Starch hydrolysis	+	-	-
Color	+	+	+

*+: Positive reaction, -: Negative reaction,

NR: Not reported

Considering all the above results, BC74171^T strain of the present invention was confirmed to be a novel microorganism belonging to *Paracoccus* genus. The present inventors named the BC74171^T strain as '*Paracoccus haeundaensis*' and deposited it at KCCM (Korean Culture Center of Microorganisms), on January 24, 2003 (Accession No: KCCM-10460). Systematic position of the *Paracoccus haeundaensis* of the present invention was shown in FIG. 6. Phylogenetic dendrogram was made out by using Treeview program. Bootstrap analysis (1000 replications) was performed using a method measuring distance and parsimony (Agnes Groisillier and Aline Lonvaud-Funel, *International Journal of Systematic Bacteriology*, 49: 1417-1428, 1999).

Example 6: Astaxanthine production in the strain
of the present invention

The strain of the present invention was cultured in 50 ml of medium (yeast extract 1%, tryptone 0.5%, NaCl 3%) at 25°C for 6 days. The cells were collected by centrifugation (13,000 rpm). The collected cells were suspended in 20 ml of methanol, which were then cultured overnight at 4°C. Centrifugation was performed again with 13,000 rpm to obtain supernatant. Optical density was measured at 190-900 nm.

As a result, as shown in FIG. 7, peaks were observed both at 450 nm and at 470 nm, and those peaks were confirmed to be original peaks of β -carotene and astaxanthine.

For the accuracy of analysis, 1 ml of the supernatant was taken and filtered by a 0.45 μ m filter, followed by HPLC analysis (column: 4.6 \times 250 mm, uBondapak C18, Waters, Milford, MA; mobile phase: acetonitrile-methanol-water(49:44:7 v/v), Flow: 10 ml/min, Detector: 470 nm). β -carotene and astaxanthine purchased from Sigma were used as a standard substance. As a result, the strain of the present invention did produce astaxanthine and

β -carotene. The amount of astaxanthine produced from the strain was 25 $\mu\text{g/g}$ (dry weight). Therefore, it was confirmed that bright orange color of the strain was caused by carotenoid biosynthesis in a cell and a major pigment accumulated in the cell was astaxanthine.

Example 7: Preparation of genomic DNA for the cloning of a carotenoid biosynthesis gene

10 *Paracoccus haeundaensis* (KCCM-10460) was cultured in a PPES-II medium (tripton 1 g/l, bacto-soyton 1 g/l, ferric citrate 0.01 g/l, polypepton 2 g/l and NaCl 3 g/l) at 25°C for 10 days. Then, the culture solution was centrifuged at 13,000 rpm to collect cells. In order to isolate genomic DNA from cells, the cells were suspended in STE buffer solution (10 mM Tris, 1 mM EDTA, 100 mM NaCl, pH 8.0), which was further reacted at 68°C for 15 minutes. The cells obtained from centrifugation were re-suspended in solution I (50 mM glucose, 25 mM Tris, 10 mM EDTA, pH 8.0). 5 mg/ml of lysozyme and 100 $\mu\text{g/ml}$ of RNase A were added, followed by reaction at 37°C for 1 hour. Then, 250 $\mu\text{g/ml}$ of proteinase K was added,

followed by reaction at 37°C for 3 hours. N-lauroylsarcosine was added by 1% of total volume, followed by reaction at 37°C. Genomic DNA was purified by phenol-chloroform extraction method.

5 Extraction was performed by adding same volume of phenol-chloroform and genomic DNA was precipitated by adding 100% ethanol twice as much as the whole volume. The precipitated DNA was washed with 70% ethanol. TE buffer solution was added to dissolve

10 the precipitated DNA completely at 65°C, which was then ready to be used.

Example 8: Construction of genomic DNA library

<8-1> Preparation of a cosmid vector

15 Xba I (9 U/ μ g) restriction enzyme was added to 25 μ g of a cosmid vector (SuperCos 1 Cosmid Vector, Stratagene), making the total volume of reacting solution 200 μ l, and the solution was reacted at 37°C for 1 hour, followed by digestion.

20 Vector DNA was purified by phenol-chloroform extraction method, which was precipitated by 100% ethanol. For dephosphorylation of the vector digested with Xba I, CIAP enzyme (Promega) was added, followed by reaction at 37°C for 30 minutes.

Then, the purified vector was reacted with *Bam*H I (5 U/ μ g) at 37°C for 1 hour. Extraction was performed by phenol-chloroform extraction method, and precipitation was induced by using ethanol. 5 The precipitate was dissolved in TE buffer solution, making the concentration 1 μ g/ μ l.

<8-2> Construction of genomic DNA library

100 μ g of *Paracoccus haeundaensis* genomic DNA 10 obtained in the above Example 7 was treated with *Sau*3A I (10 U), inducing a partial enzyme reaction. Upon completing the enzyme reaction, 0.5 M EDTA was added. Then, genomic DNA was separated by phenol-chloroform extraction method, which was 15 precipitated by 100% ethanol. Genomic DNA, a product of a partial enzyme reaction, was dissolved in TE buffer solution, and treated with *CIAP* enzyme by the same method as used in the above Example 7, leading to dephosphorylation. 20 DNA was separated again by phenol-chloroform extraction method. For the ligation with a cosmid vector prepared in the above Example <8-1>, T4 ligase (Promega) and 10x ligase buffer (Promega) were added to the isolated DNA, followed by 25 reaction at 12°C for 18 hours. After completing

the reaction, an *E. coli* strain XL1-Blue (Stratagene) was transformed with the ligation mixture, resulting in the construction of a genomic DNA library.

5

Example 9: Examination and analysis of a transgenic strain containing a color-producing gene

FPP (Sigma), one of common substrates included in carotenoids, was added to a LB agar medium by 1% out of total volume. A genomic library constructed in the above Example 8 was smeared on the plate, which was cultured at 37°C. Among cultured colonies, the ones having orange color were selected (13 out of about 2000 colonies). Cosmid vectors were isolated from those 13 colonies selected above. Then, primer working sequencing was performed to identify nucleotide sequences of DNA fragments inserted in each cosmid vector. Identification of the nucleotide sequences was committed to GenoTech Corp., Korea.

As a result, the smallest DNA insert of all fragments inserted in the cosmid vectors was 6,223 bp long, and had nucleotide sequences represented by SEQ. ID. No 4. A cosmid vector containing the
5 nucleotide sequence represented by SEQ. ID. No 4 was named 'COSCRT'.

Example 10: Sequence analysis of a DNA insert containing a carotenoid biosynthesis related gene

10 Sequence analysis of a DNA insert obtained in the above Example 9 was performed using NCBI ORF Finder program
(<http://www.ncbi.nlm.nih.gov/gorf/gorf.html>) to analyze ORF.

15 As a result, 6 ORFs were included in a DNA insert obtained in the above Example 3. Each ORF showed a relevant homology with nucleotide sequences of *crtW* coding β -carotene ketolase, *crtZ* coding β -carotene hydroxylase, *crtY* coding
20 lycopene cyclase, *crtI* coding phytoene desaturase, *crtB* coding phytoene synthase and *crtE* coding geranylgeranyl pyrophosphate synthase. All of those enzymes were confirmed to be involved in carotenoid biosynthesis.

The comparison of homology between amino acid sequences translated from each ORF and amino acid sequences of each carotenoid biosynthesis enzyme isolated from *Alcaligenes* sp. (Misawa et al., 5 *Biochem. Biophys. Res. Commun.*, 209(3): 867-876, 1995), *Bradyrhizobium* sp. (Hannibal et al., *J. Bacteriol.*, 182(13): 3850-3853, 2000) and *Flavobacterium* sp. (Pasamotes et al., *Gene*, 185(1): 35-41, 1997) was shown in FIG. 9 and FIG. 10 14. The nucleotide sequences of 6 ORF genes cloned by the present inventors and amino acid sequences translated therefrom were each represented by SEQ. ID. No 5 ~ No 16. In particular, *crtW*, *crtZ*, *crtY*, *crtI* and *crtB* genes 15 were each represented by SEQ. ID. No 5, No 7, No 9, No 11, No 13 and No 15 and amino acid sequences of each gene above were represented by SEQ. ID. No 6, No 8, No 9, No 12, No 13 and No 16, respectively.

It was confirmed from the above results that 20 *crt* gene involved in biosynthesis of carotenoid was included in a DNA fragment inserted in the cosmid vector above.

The composition of *crt* gene of the present invention was shown in FIG. 15. As shown in FIG. 25 15, termination codon and start codon were

overlapped in each *crtW*, *crtZ*, *crtY*, and *crtI*.
Especially, *crtE* gene seemed to have a
directionality of complementary strands, and had
each of recognition sequences of *KpnI*, *XmaI*, *SmaI*,
5 *ClaI*, *HindIII* and *BamH* in its sequences.

Example 11: Expression of *crt* gene in *E. coli*

In order to investigate if carotenoid could
be produced by a protein expressed by *crt* gene of
10 *Paracoccus haeundaensis* isolated in the above
Example 9, *crt* gene was first amplified by PCR
using HL premix (Bioneer). At that time,
oligoneucleotide primers represented by SEQ. ID.
No 17 and No 18 were used. All the PCRs were
15 performed as follows; pre-denaturation at 94°C for
5 minutes, denaturation at 94°C for 30 seconds,
annealing at 66°C for 30 seconds, polymerization
at 72°C for 6 minute, 25 cycles from denaturation
to polymerization, and final extension at 72°C for
20 20 minutes. The PCR product was inserted in Topo-
XL-vector (Invitrogen), which was used for
transduction of *E. coli*. At that time, XL1-Blue
(Stratagene), TOPO (Invitrogen), BL21(DE3) codon
plus (Stratagene), DH1 (Takara) and DH5a (Takara)

were used as *E. coli* of the present invention. As a result, all transformants of BL21(DE3), XL1-Blue, BL21(DE3) codon plus were confirmed to have orange color. Thereafter, each transformant of *E. coli* was selected for culture. As a result, transformed BL21(DE3) codon plus produced astaxanthine most.

The present inventors inserted *crt* gene having nucleotide sequences represented by SEQ. ID. No 4, isolated in the above Example 9, into pCR-XL-TOPO vector (Invitrogen), a gene expression vector, which was then named 'pCR-XL-TOPO-crtfull' (FIG. 16). BL21(DE3) codon plus cells were transfected with the vector 'pCR-XL-TOPO-crtfull', followed by selection of strains having orange color. The selected strains were cultured in 50 ml of LB medium at 37°C for 8 hours. The culture solution was centrifuged at 13,000 rpm, then supernatant was discarded and cells were collected. 20 ml of methanol was added to the collected cells. After vortexing, the cells were cultured at 4°C for overnight. Centrifugation was performed (13,000 rpm) to obtain supernatant. In order to confirm whether carotenoid was generated, optical

density was measured at 190-900 nm and 400-550 nm. As a result, peaks were observed at 450 nm and 470 nm, which were confirmed to be original peaks of β -carotene and astaxanthine (FIG. 17).

5 For the accuracy of analysis, 1 ml of the above supernatant was obtained and filtered with a 0.45 μ m (pore size) filter, followed by HPLC analysis (column: 4.6x250mm, uBondapak C18, Waters, Milford, MA; mobile phase: acetonitrile-methanol-water(49:44:7 v/v), Flow: 10ml/min, Detector: 10 470nm). At that time, β -carotene and astaxanthine, purchased from Sigma, were used as a standard substance. As a result, the substances produced from the strain of the present invention 15 were confirmed to be astaxanthine and β -carotene.

Astaxanthine produced from the strain of the present invention was also quantified, resulting in the production of 110 μ g/g (dry weight). For that result, neither inducer nor carotenoid 20 substrate was added, suggesting that β -carotene and astaxanthine could be produced in *E. coli* only by nucleotide sequences having 6,223 bp isolated in the present invention. The amount of astaxanthine quantified above was far more than 25 that produced (25 μ g/g) by *Paracoccus haeundaensis*

(Accession No: KCCM-10460).

INDUSTRIAL APPLICABILITY

As explained hereinbefore, the present
5 inventors have isolated and identified a novel
Paracoccus genus microorganism mostly producing
astaxanthine among many carotenoids, and also have
cloned 6 genes coding proteins involved in
carotenoid biosynthesis and *crt* gene containing
10 the same from the above microorganism. The
present inventors also have confirmed that
carotenoid can be produced even in *E. coli* not
producing carotenoid, by using *crt* gene.
Therefore, a gene of the present invention and a
15 microorganism producing the gene can be
effectively used for the production of carotenoids
such as β -carotene and astaxanthine which are
available for making food, medicines and beauty
stuffs.

20

Those skilled in the art will appreciate that
the conceptions and specific embodiments disclosed
in the foregoing description may be readily

utilized as a basis for modifying or designing other embodiments for carrying out the same purposes of the present invention. Those skilled in the art will also appreciate that such

5 equivalent embodiments do not depart from the spirit and scope of the invention as set forth in the appended claims.

What is claimed is

1. A gene coding a protein involved in carotenoid biosynthesis, which has nucleotide sequences
5 selected from a group consisting of nucleotide sequences represented by SEQ. ID. No 5, No 7, No 9, No 11, No 13 and No 15.
2. The gene as set forth in claim 1, wherein the
10 gene has nucleotide sequences of *crtW* coding β -carotene ketolase and represented by SEQ. ID. No 5.
3. The gene as set forth in claim 1, wherein the
15 gene has nucleotide sequences of *crtZ* coding β -carotene hydroxylase and represented by SEQ. ID. No 7.
4. The gene as set forth in claim 1, wherein the
20 gene has nucleotide sequences of *crtY* coding lycopene cyclase and represented by SEQ. ID. No 9.
5. The gene as set forth in claim 1, wherein the
25 gene has nucleotide sequences of *crtI* coding

phytoene desaturase and represented by SEQ. ID.
No 11.

5 6. The gene as set forth in claim 1, wherein the
gene has nucleotide sequences of *crtB* coding
phytoene synthase and represented by SEQ. ID.
No 13.

10 7. The gene as set forth in claim 1, wherein the
gene has nucleotide sequences of *crtE* coding
geranylgeranyl pyrophosphate synthase and
represented by SEQ. ID. No 15.

15 8. A *crt* gene containing all the genes of claim 2
~ claim 7 and represented by SEQ. ID. No 4.

20 9. A protein encoded by the gene of claim 1,
which has nucleotide sequences selected from a
group consisting of nucleotide sequences
represented by SEQ. ID. No 6, No 8, No 10, No
12, No 14 and No 16.

25 10. A recombinant vector containing the *crt* gene
of claim 8.


11. The recombinant vector as set forth in claim 10, wherein the vector is pCR-XL-TOPO-crtfull having a cleavage map represented in FIG. 16.
- 5 12. An *E. coli* transformant transformed with the recombinant vector of claim 11.
13. A method for producing carotenoid comprising the following steps:
- 10 1) Cloning the *crt* gene of claim 8;
- 2) Constructing a recombinant vector in which the *crt* gene of the above step 1) was inserted;
- 15 3) Transfecting a host cell with the recombinant vector of the step 2); and
- 4) Recovering carotenoids from the culture cells in which a strain transformed with the above recombinant vector was being cultured.
- 20 14. The method as set forth in claim 13, wherein the recombinant vector is that of claim 11.
- 25 15. The method as set forth in claim 13, wherein the host cell is *E. coli* or yeast.

16. The method as set forth in claim 13, wherein
the recovery of carotenoids is performed from
the culture cells in which the *E. coli* was
5 being cultured.
17. The method as set forth in claim 13, wherein
the cartenoid is β -carotene or astaxanthine.
- 10 18. A *Paracoccus haeundaensis* producing
astaxanthine, which has a 16S rDNA nucleotide
sequence represented by SEQ. ID. No 3.
- 15 19. The *Paracoccus haeundaensis* as set forth in
claim 18, wherein the strain is represented by
accession No: KCCM-10460.

1/11

FIGURES

FIG. 1



2/11

FIG. 2





FIG. 3



3/11

FIG. 4






FIG. 5




4/11

FIG. 6




5/11

FIG. 7



6/11

FIG. 8



7/11

FIG. 9

	1		100
<i>P. haeundaesis</i>	(1)	NSAHLEPKADLTATSLIYSGGIIAAVLALHVALWFLDAAAHPLAIANEIGLT-WLSYGLPFIABDAHBGSVYIGRPBMAAMQQLVLR	
<i>Alcaligenes_sp</i>	(1)	NSGRKPGTIGDTIVNELGLTAAILLCPVLHAFILWLLDAAAHPLCAVCLAGLT-WLSYGLPFIABDAHBGSVYIGRPBMAAMQQLALW	
<i>Bradyrhizobium_sp</i>	(1)	MHAATANATEFGASRRDDABORRVGLTAAVITAAALVHVGLENFEPLTLHSLLPALPLVYVLTWLYKLEFIABDCBRSIYPPEPQVHBRIQQLCLP	
Consensus	(1)	NSA K A T V L LSAALIAAALVLRV LWFDAAAHPLLAII LGLY WLSYGLPFIABDAHBGSVYIGRPBMAAMQQL LV	
	101		200
<i>P. haeundaesis</i>	(90)	LYAGFSVRKNIYKHNHHRBTGTDDEDFDHC--GPVRVYARFIGTYEGWRBGLLPLVITVYALILGD-RWHYVFWPLPSILASIQLEFVFWLPHR	
<i>Alcaligenes_sp</i>	(90)	LYAGFSVRKLIYKHNHHRBTGTDDEDFDHC--GPVRVYGSFVSTYEGWRBGLLPLVITVYALILGD-RWHYVFWPVPVAVLASIQLEFVFWLPHR	
<i>Bradyrhizobium_sp</i>	(101)	LYAGFSFDALNVHRHHRBPGTAEDDFEDVPPKGFVWHPASFELHYFGKQVATIAVSLVYQLVFAVPLQWILLFQALPGLISADQLETPTTYLRK	
Consensus	(101)	LYAGFSV KLIYKHN HRRH GTDDDFDHC GPVRVYASFI TYEGWRBGLLPLVITVYALILGD RWHYVFWPLPAILASIQLEFVFWLPHR	
	201		259
<i>P. haeundaesis</i>	(186)	PGHDAPFDRHNAISSRISDPVSLLCFHFEGGYEHEHLEPTVPVWRLPSTRTKGDTA--	
<i>Alcaligenes_sp</i>	(186)	PGHDDFDRHNAISTGIGDPLSLTCFHFEGGYEHEHLEPTVPVWRLPETRTKTCGBA--	
<i>Bradyrhizobium_sp</i>	(201)	PATQPFADHNAISTSEPAVLSLLTCFHFEGGYEHEHLEPTDAPQVRLPEIKRBALERRD	
Consensus	(201)	PGHD FDRHNAISS I DPLSLLCFHFEGGYEHEHLEPTVPVWRLP TRKGD A	

FIG. 10

	1		100
<i>P. haeundaesis</i>	(1)	HTNFLIVVATVLVHETAYSVRRGTHRPLGQGGKSHHEERDHALEKNDLYGLVFAVIATVLT VQWIDAPVLWJIALGHTVGLIYFVLHDLVHQRW	
<i>Alcaligenes_sp</i>	(1)	HTQPLIVVATVLVHETAYSVRRGTHRPLGQGGKSHHEERDHALEKNDLYGLVFAVLATILET VQAYWQPVLVJIALGHTVGLIYFVLHDLVHQRW	
Consensus	(1)	HTNFLIVVATVLVHETAYSVRRGTHRPLGQGGKSHHEERDHALEKNDLYGLVFAVIATVLT VQV W PVLWJIALGHTVGLIYFVLHDLVHQRW	
	101		163
<i>P. haeundaesis</i>	(101)	PFRYIPRKYABLYQAHLEHHAVEGRDHCVSFGFIYAPPYDKLKQDLKTSGVLEBAHQERT	
<i>Alcaligenes_sp</i>	(101)	PFRYIPRKYABLYQAHLEHHAVEGRDHCVSFGFIYAPPYDKLKQDLKTSGVLEPQDERPS	
Consensus	(101)	PFRYIPRKY RRLYQAHLEHHAVEGRDHCVSFGFIYAPPYDKLKQDLK SGVLB S	

8/11

FIG. 11

	1	100
P. haemundae	(1) VTRDVLGAGLGLALALBAERDLVLLDHAQPSDGHTECHDPLSPHVLAKLPERRANWPDQEVFPBABELATGYSLDGAALADAYAB	
Flavobacterium_sp	(1) MSBDLLIAGAGLSGALIALAVEDRETDARI VMLUASSQPSIQHTCCHDTDLSPVLAKLPSPLESEYDQFAFPDESELTGYGSIAGALIGLLQ-	
Consensus	(1) MSBDLLIAGAGLA ALIALALR RPD RI LLLD AQPSD HTSCSD DLSP VLAKLP PIRRA R DQEV FP BAREL TGYGSIIDAAAL L	
	101	200
P. haemundae	(101) SGAETDWSIDIALDEQATLSGTRIAGAVLDGGAQPSHLTVGFQEPVGVKIEYDCFPQWPPHEDATVTQDDGYEPIILLPFSTRILIEDYBY	
Flavobacterium_sp	(100) -GVDLSWHTVATLDOTGATLTDSSELEAACVIDALQAVETPLTVPQKTPGVKIEYDAPHGVVERPHEDATVPQDDVTFVYLLPFSTRILIEDYBY	
Consensus	(101) G DIBWS IA LDD GATLS GSRLEAA VIDARGA S HLTVGFQEPVGVKIEYD PHGV BPHINDATV Q DGYEPIYLLPFSTRILIEDYBY	
	201	300
P. haemundae	(201) SDGSLDDALAAERDYARQQGVIGARVREDGILEYALAHDAAPFVADHESGPVPGVLRAGEFHPVTGYLLTAQVADVVAGLSGPPGTDALEGAIP	
Flavobacterium_sp	(189) SUGLDLGGALAAQLDYAARQGVIGQEMKREMGITPIALAHDAIGFVDRHAGQAVPPVGLGALHPTVTGSEPTAAQVADAIAR--DLTTASABRAYE	
Consensus	(201) SDGG LDD ALA AS DYA GWTG EMREBERGILPITALANDA GFW DBA G VPPGL AG FHPVTGYSLPTAAQVAD IAA T A R AIR	
	301	387
P. haemundae	(301) DYADEAREDEFILRLRLPRGAPDRBTLLQEPVNPESLI ESYAGELSVADQLRI VTKPPTPLGTATDCLPERPLKENA	
Flavobacterium_sp	(297) GWALLADMSRFLLEMLHLPCTPPDRBTLLQEPVNPESLI ESYAGELSVADQLRI VTKPPTPLGTATDCLPERPLKQBR	
Consensus	(301) VADRA DRFLRLRLNRLPRGCP DERY LLQRFYRLP LIERFYAGELSLAD LRIVTKPPIPL ALRCLPERPL E A	

FIG. 12

	1	100
P. haemundae	(1) HNAESPAANTKIVIGGEGALALALRQSGAGIATVLEADDEGCHATTSHRQGVFEDAGPTVTIDPDALKELALTOGDHARDVTLHVCTFFELHWP	
Flavobacterium_sp	(1) -----HSSATVIGAGPQGLALALRQSGAGIATTVBARHKGGRAYVVMHQGVFEDAGPTVTIDPDALKELALTOGDHARDVTLHVCTFFELHWP	
Consensus	(1) S TVIGAGPQGLALALRQSGAGIATTVBARHKGGRAYVV DQGVFEDAGPTVTIDPDALKELALTOGDHARDVTLHVCTFFELHWP	
	101	200
P. haemundae	(101) GEFVDTFHEALQELQIAQFLDLEGYRSEFDYNEETQDQVMAQVFLGGLVLRKAPALAKLEAYEC SHAKWATITKPEVLLDANSTELLLWQH	
Flavobacterium_sp	(94) GRSFEYVDDDELELVVSEFTQADVDTSEHDFVSEVYVRLQELATTCPULKGLQENNAAPALAKLEAYEC VHSHPARFQDEHELPACTFEILLWQH	
Consensus	(101) GR FDYND D L RQIA FNP DLQGVRR DYAREVY EGVLEKLT PFLKLGQL AAPALHKL AYKSVHA VA FI DPHLRQAFSPHLLWQH	
	201	300
P. haemundae	(201) PFSTSSITALIBALERGGVFPAGOSTRQVAGHVALFERLGQNNLHAKVARTETDQARTCYVLADGESLEADNVAAGVHNVBLGLETAGQSR	
Flavobacterium_sp	(194) PFSTSSITALIBALERGGVFPAGOSTHOLVAGHVALFERLGQ LLLHARV RIDEGR B FGVTL DGR LEAD VASNGDVH YEDLGHT RG SK	
Consensus	(201) PFSTSSITALIBALERGGVFPAGOSTHOLVAGHVALFERLGQ LLLHARV RIDEGR B FGVTL DGR LEAD VASNGDVH YEDLGHT RG SK	
	301	400
P. haemundae	(301) AKSLDEKRVKSLFVLFHGLREAPEDIAEHTILFQPVYRELVHEIFEGFLAEDFSLYLESPECTDQDAPPGESTBYVLPAPVFLQDABIDHAVRGVRY	
Flavobacterium_sp	(294) AAILNRQVCKSLFVLFHGLSKNFENLAHSHVIFQPEFEGLVREIFNUPRLDDFSHYLHSPVYVTPSLAIEGHSTHYVLPAPVFLQDABIDHAVRGVRY	
Consensus	(301) A L R RVNSILFVLFHGL P IAHSHIFGPVRY LVNEIF GPKL DDFSLYLHSPC TDP LAP GHSTHYVLPAPVFLQDABIDHAVRGVRY	
	401	500
P. haemundae	(401) ADRIASLERELIPNLRANLITTRIPADFASLHNADEGAFSVEPIITQCAVFRPHNRDXTIRNFFLVGAGTHPGAGIPGVGSAKATAQVHLSDLA	
Flavobacterium_sp	(394) AERIIFEREREALPDLRKLTVSEIFSPADESTELS AHHGSAFVEPIITQCAVFRPHNRDXTIRNFFLVGAGTHPGAGIPGVGSAKATAQVHLSDLA	
Consensus	(401) ADRI LE R IP LR LT SRIFSPADFASL AHHGSAFVEPIITQCAVFRPHNRDXTIRNFFLVGAGTHPGAGIPGVGSAKATAQVHLSDLA	
	501	
P. haemundae	(501) A	

9/11

FIG. 13

1 100

P. haemdaesis (1) HSDLVLTSTETATQSSQSFATAAKLHPGIRDDTVHLYANCRHADDYIDQALQSRPEAVNDPQARLDGLSDVTLAALQGGPVTFPAALDAYASSRDF
 Flavobacterium_sp (1) HTDLTATSEAAI AQGSCSFAQAARLPPGIRDDTVHLYANCRHADDYIDQVHSSAPRAGGTPQARLGALRAVTLAALHEDGPNSPFFAALQVABRDF
 Consensus (1) HSDL TS AI QGSCSFA AAKLHPPGIRDDTVHLYANCRHADDYIDQ LGS PBA DPQARL ALR DTLAAL DGPNSPPFAALR VARBRDF

101 200

P. haemdaesis (101) PQAWPHDLIEGFANDVEARDVRLDDVLEYSTYVAGIVGTTHARVGVYRDPVLDRACDLGLAFCLTHIARDVIDDARIGCYLPGVDLDQAGARIDGPF
 Flavobacterium_sp (101) EDLVPDLIEGFANDVADEYRSLDDVLEYSTYVAGIVGTTHARVGVYRDPVLDRACDLGLAFCLTHIARDVIDDAAIGCYLPGVDLDQAGARIDGPF
 Consensus (101) P WENDLIEGFANDV EDYRSLDDVLEYSTYVAGIVGTTHARVGVY DD VLDRACDLGLAFCLTHIARDVIDDA IGRCYLPGVDL AGA IDGPF

201 300

P. haemdaesis (201) ESPELTYVILRLDRAEYTYASAKVGLADLPFCASIAAALRI THAI GLRENSGPQATQRTSKSKAAKIGLLSYGGVDVARSRLPGAGVSRGGQLYE
 Flavobacterium_sp (201) ECDALYSVILRLDRAEYTYASAKVGLADLPFCASIAAALRI THAI GLRENSGPQATQRTSKSKAAKIGLLSYGGVDVARSRLPGAGVSRGGQLYE
 Consensus (201) ES LYSVILRLD AEPYTYASR GL LPFCASIAAALRI THAI RIR GP AYEQRISTSKAAKIGLLA GG D A SRL GA ISR GLWTR

301

P. haemdaesis (301) PHEV
 Flavobacterium_sp (301) PBA
 Consensus (301) P

FIG. 14

1 100

P. haemdaesis (1) HRRDVNPIHATILQTKLEETAGGTCAVSQILQANRCHGMLSSRRFLGHLLELLAAEASGGGCTIYDAACAPENHNASLIDPDLCTDAGLASGRCP
 Flavobacterium_sp (1) HTPQPPPLRLDVEIRLAQISGGQVTVSAPLQAAEDDAALSPKLFRAVLHLNVAESSQVCDANVDAACAVENHNASLIDPDLCTDAGLASGRCP
 Consensus (1) N N LL RL IA FGVS PLGAANS AALS GKRFAHLMLL ABAAGGVCD IVDAACAVENHNASLIDPDLCHDDA BRG PAT

101 200

P. haemdaesis (101) HWAGSEDAVIGQIALITYAHALLAGAGASCTVRAQLVRLSSSLAQGLCGQDLRLHAAKNGAGVQEQIDLYTGVLTAGLEHLAVIEFPAEDDQIQ
 Flavobacterium_sp (101) HWAGSEDAVIGQIALITYAHALLAGAGASCTVRAQLVRLSSSLAQGLCGQDLRLHAAKNGAGVQEQIDLYTGVLTAGLEHLAVIEFPAEDDQIQ
 Consensus (101) HWAGSEDAVIGQIALITYAHALLAGAGASCTVRAQLVRLSSSLAQGLCGQDLRLHAAKNGAGVQEQIDLYTGVLTAGLEHLAVIEFPAEDDQIQ

201 295

P. haemdaesis (201) HIDEFGLGRVFGSYDLDLVYVGHQALGLITGBDAAPGPRWILLAYSIDLQNVSRHYEASRAQLPAMLSKRLQAPRIAAALERTVPTAALA--
 Flavobacterium_sp (201) LHAPEGLGRVFGSYDLDLVYVGHQALGLITGBDAAPGPRWILLAYSIDLQNVSRHYEASRAQLPAMLSKRLQAPRIAAALERTVPTAALA--
 Consensus (201) LI FGRGLGRVFGSYDLDLVYVGHQALGLITGBDAAPGPRWILLAYSIDLQNVSRHYEASRAQLPAMLSKRLQAPRIAAALERTVPTAALA--

10/11

FIG. 15







FIG. 16



11/11

FIG. 17



SEQUENCE LISTING

<110> KIM, Young Tae
 LEE, Jae Hyung
 ALGENETECH

<120> Gene involved in the biosynthesis of carotenoid and marine
 microorganism, paracoccus haeundaesis, producing the
 carotenoid

<130> 4fpo-02-06

<150> KR2003-20222
 <151> 2003-03-31

<150> KR2003-20023
 <151> 2003-03-31

<160> 18

<170> KopatentIn 1.71

<210> 1
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> forward primer for Paracoccus haeundaesis 16S rDNA

<400> 1
 cataagtaat tatggttttg t 21

<210> 2
 <211> 18
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> reverse primer for Paracoccus haeundaesis 16S rDNA

<400> 2
 cgcttcctta gaaaggag 18

<210> 3
 <211> 1454
 <212> DNA
 <213> Paracoccus haeundaesis

<400> 3
 caacttgaga gtttgatoc t ggctcagaac gaacgctggc ggcaggctta acacatgcaa 60

```

gtcgcgagcgag accttcgggt ctagcggcgg acgggtgagt aacgcgtggg aacgtgccct 120
tctctacgga atagccccgg gaaactggga gtaataccgt atacgccctt tgggggaaag 180
atztatcgga gaaggatcgg ccgcggttg attaggtagt tgggtgggta atggcccacc 240
aagccgacga tccatagctg gtttgagagg atgatcagcc aactggggac tgagacacgg 300
cccagactcc tacgggaggc agcagtgggg aatcttagac aatgggggca accctgatct 360
agccatgccg cgtgagtgat gaaggcetta gggttgtaaa gctctttcag ctgggaagat 420
aatgacggta ccagcagaag aagccccggc taactccgtg ccagcagccg cggtaatacg 480
gaggggggcta gcgttggtcg gaattactgg gcgtaaagcg cacgtgggcg gactggaaag 540
tcagagggtga aatcccaggg ctcaaccttg gaactgcctt tgaaactatc agtctggagt 600
tcgagagagg tgagtggaat tccgagtgtg gaggtgaaat tcgtagatat tcggaggaac 660
accagtggcg aaggcggctc actggctcga tactgacgct gaggtgcgaa agcgtgggga 720
gcaaacagga ttagataccc tggtagtcca cgccgtaaac gatgaatgcc agacgtcggc 780
aagcatgctt gtcggtgtca cacctaacgg attaagcatt ccgcctgggg agtacggtcg 840
caagattaaa actcaaagga attgacgggg gccgcacaa gcggtggagc atgtggttta 900
attcgaagca acgcgcagaa ccttaccaac ccttgacatg gcaggaccgc tggagagatt 960
cagctttctc gtaagagacc tgcacacagg tgctgcatgg ctgtcgtcag ctgctgtcgt 1020
gagatgttcg gttaagtccg gcaacgagcg caaccacgt ccctagttgc cagcattcag 1080
ttgggcactc tatggaaaact gccgatgata agtcggagga aggtgtggat gacgtcaagt 1140
tctcatggcc cttacgggtt gggctacaca cgtgctacaa tgggtggtgac agtgggttaa 1200
tccccaaaag ccatctcagt tcggattgtc ctctgcaact cgagggcagt aagttggaat 1260
cgctagtaat cgcggaacag catgccgcgg tgaatacgtt cccgggcctt gtacacaccg 1320
cccgtcacac catgggagtt gggtctaccc gacgacgtg cgctaaccct cggggggcag 1380
gcggccacgg taggatcagc gactggggtg aagtcgtaac aaggtagccg taggggaacc 1440
tgcggctgga tcac 1454

```

```

<210> 4
<211> 6223
<212> DNA
<213> crt gene

```

```

<400> 4
gttccacgac tggggcatcc ccacgaccgc gtcgctgcgc gccatcgcgc cgatgatggg 60

```

gccggaccgg gttctggtcg ggtcgggagg ggtgcgtcac gggctggacg ccgcgcgggc 120
 catccgcctc ggcgcggacc tcgtggggca ggcggcccgc gcgtgcccgc ccgcgcgcca 180
 cagcgccgag gccctgtccg atcacctgtc cgacgtcgtg acccagctgc gcatcgcgat 240
 gttctgcacc ggatcggggc accttgacgc gctgcgtgc gcgcctctgc tgggtgccggg 300
 gccgggtggc caatggtcgc aagcaacggg gatggaaacc ggcgatgcgg gactgtagtc 360
 tgcgcggatc gccggtccgg gggacaagat gagcgcacat gccctgccc aaggcagatct 420
 gaccgccacc agcctgatcg tctcggggcg catcatcgcc gcgtggctgg ccctgcatgt 480
 gcatgcgctg tggtttctgg acgcggcggc gcatcccatc ctggcgatcg cgaatttcct 540
 ggggctgacc tggctgtcgg tcgggtctgt cttcatcgcg catgacgcga tgcacgggtc 600
 ggtcgtgccg gggcgctccg gcggcaatgc ggcgatgggc cagctggtcc tgtggctgta 660
 tgccggattt tcgtggcgca agatgatcgt caagcacatg gccatcacc gccataccgg 720
 aaccgacgac gaccccgatt tcgacctagg cggcccggtc cgttggtacg cgcgcttcat 780
 cggcacctat ttcggctggc gcgaggggct gctgctgcc gtcacgtga cgggtctatgc 840
 gctgacctg ggggatcgt ggatgtacgt ggtcttctgg ccgctgccgt cgatcctggc 900
 gtcatccag ctgttcgtgt tcggcacctg gctgccgcac cggcccgcc acgacgcgtt 960
 cccggaccgc cataatgcgc ggtcgtcgc gatcagcgac ccgctgtcgc tgctgacctg 1020
 ctttcacttt ggtggttatc atcacgaaca ccacctgcac ccgacggtgc cttggtggcg 1080
 cctgcccagc acccgacca agggggacac cgcatgacca atttcctgat cgtcgtcgcc 1140
 accgtgctgg tgatggagtt gacggcctat tccgtccacc gttggatcat gcacggcccc 1200
 ctgggctggg gctggcaca gtcccaccac gaggaacacg accacgcgct ggaaaagaac 1260
 gacctgtacg gcctggtctt tcgggtgatc gccacggtgc tgttcacggt gggctggatc 1320
 tgggcgcggg tcctgtggtg gatcgctttg ggcgatgccg tctatgggct gatctatttc 1380
 gtcctgcatg acgggctggt tcatcagcgc tggccgttcc gctatatccc gcgcaagggc 1440
 tatgcccgc gcctgtatca ggcccaccgc ctgcaccacg cggtcgaggg acgcgacct 1500
 tgcgtcagct tcggcttcat ctatgcgcc cgggtcgaca agctgaagca ggacctgaag 1560
 acgtcggggc tgctgcgggc cgaggcgag gagcgcacgt gacctatgac gtgctgctgg 1620
 caggggcggg ccttgcaac gggctgatcg ccctggcgct gcgcgcggcg cggcccgcacc 1680
 tgcgggtgct gctgctggat catgcggcg gaccgtcaga cggccatacc tggctcctgcc 1740
 acgacccga tctgtcgcc cactgggtgg cgcgggtgaa gccctgcgc cgcgccaact 1800

ggcccgacca ggaggtgcgc tttccccgcc atgcccggcg gctggccacc ggttacgggt 1860
cgctggacgg ggcggcgctg gcggatgcgg tggccccgtc gggcgccgag atccgctgga 1920
acagcgacat cgccctgctg gatgaacagg gggcgacgct gtcctgcggc acccggatcg 1980
aggcggggcg ggtcctggac gggcgcggcg cgcagccgtc gcggcatctg accgtgggtt 2040
tccagaaatt cgtgggcgtc gagatcgaga ccgactgcc ccacggcgtg ccccgcccga 2100
tgatcatgga cgcgaccgtc acccagcagg acgggtaccg attcatctat ctgctgccct 2160
tctctccgac gcgcatcctg atcgaggaca ctcgctatc cgatggcggc aatctggacg 2220
acgacgcgct ggcggcggcg tcccacgact atgcccgcca gcagggctgg accggggccg 2280
aggtcggcg cgaacgcggc atcctgccc a ttgcgtggc ccatgacgcg gcgggcttct 2340
gggccgatca cgcggagggg cctgttccc tgggactgcg cgcgggggtt tttcaccgg 2400
tcaccggcta ttcgctgcc tatgcggcg aggtggcgga cgtggtggcg ggctgtccg 2460
ggccgcccgg caccgacgcg ctgcgcggcg ccatccgca ttacgcgatc gaccgggcac 2520
gccgtgaccg ctttctgcgc ctgctgaacc ggatgctgtt ccgcggtgc gcgcccgacc 2580
ggcgctatac cctgctgcag cggttctacc gcatgccga tggactgatc gaacggttct 2640
atgccggccg gctgagcgtg gcggatcagc tgcgcatcgt gaccggcaag cctccattc 2700
cccttggcac ggccatccgc tgcctgccg aacgtcccct gctgaaggaa aacgcatgaa 2760
cgccattcg cccgcggcca agaccgcat cgtgatcggc gcaggctttg gcgggctggc 2820
cctggccatc cgctgcagt ccgcgggcat cgccaccacc ctggtcgagg cccgggacaa 2880
gcccggcggg cgcgcctatg tctggcacga tcagggccat gtcttcgacg cgggcccga 2940
cgtcatcacc gaccccgatg cgctcaagg gctgtggcg ctgaccggc aggacatggc 3000
gcgcgacgtg acgctgatgc cgggtgcgcc cttctatcga ctgatgtggc cggcgggaa 3060
ggtcttcgat tacgtgaacg aggcgatca gctggagcgc cagatcgccc agttcaacc 3120
ggacgacctg gaaggatacc gccgcttcg tgattacgcg gaggaggtgt atcaggagg 3180
ctacgtcaag ctgggcaccg tgccttcct caagctggc cagatgctca aggcgcgcc 3240
cgcgctgatg aagctggagg cctataagtc cgtccatgcc aaggtcgca ccttcatcaa 3300
ggaccctat ctgcggcagg cgttttcgta tcacacgtg ctggtggcg ggaatccctt 3360
ctcgaccagc tcgatctatg cgctgatcca cgcgctggag cggcgcgcg gggctctggtt 3420
cgccaagggc ggcaccaacc agctggtcgc gggcatggtc gcgctgttcg aacggcttg 3480

cggccagatg atgctgaacg ccaaggctgc ccggatcgag accgagggcg cgcggaccac 3540
 gggcgtcacc ctggcggacg ggcggtcttt aagggccgac atggtcgcca gcaacggcga 3600
 cgctcatgcac aactatcgcg acctgctggg ccacacggcc cgcgggcaga gccgcgcgaa 3660
 atcgctggac cgcaagcgct ggtccatgtc gttgttcgtg ctgcatttcg gtctgcgcga 3720
 ggcgcccag gacatcgcg atcacacccat cctgttcggc ccccgctaca gggagctggt 3780
 caacgagatc ttcaagggcc cgaagctggc cgaggatttc tcgctgtacc tgcattcgcc 3840
 ctgcacgacc gatccggaca tggcgccctcc gggcatgtcc acgcattacg tgctggcccc 3900
 cgtgccgcac ctggggcccg ccgagatcga ttgggcggtc gagggggccg gctatgcga 3960
 ccgcatectg gcgtccctgg aggagcggct gatcccgaac ctgcgcgcca acctgaccac 4020
 gacgcgcac ttacgccc ccatcttcgc cagcgaactg aacgcccac acggcagcgc 4080
 cttctcggtc gagccgatcc tgacgcaatc cgcgtggttc cggccgcaca accgcgacaa 4140
 gacgatccgc aacttctatc tggtcggcgc gggcaccat cggggcgcgg gcattccggg 4200
 cgtcgtgggc tcggccaagg ccacggccca ggtgatgctg tccgacctg cgggcgcacg 4260
 agcgatctgg tcctgacctc gaccgaggcg atcacccaag ggtcgcaaag ctttgccacg 4320
 gcggccaagc tgatgccgc gggcatccgc gacgacacg tgatgctcta tgcctggtgc 4380
 cgccacgcgg atgacgtgat cgacggtcag gccctgggca gccgccccga ggcggtgaac 4440
 gaccgcagc cgcggtgga cggcctgcgc gtcgacacgc tggcgccct gcagggcgac 4500
 ggtccggtga ccccgccctt tgccgcgctg cgcgcggtgg cgcggcgga tgatttcccg 4560
 caggcctggc ccatggacct gatcgaaggc ttcgcgatgg atgtcgaggc gcgcgactat 4620
 cgcacgctgg atgacgtgct ggaatattcc tatcacgtcg caggcatcgt cggcgtgatg 4680
 atggcccgcg tgatggcggt gcgcgacgat cctgtcctgg accgcgcctg cgacctgggg 4740
 ctggcggttc agctgaccaa catcgcgcg gacgtgatcg acgatgcgc catcgggcgg 4800
 tgctatctgc cgggggactg gctggaccag gcgggcgcgc ggatcgacgg gccggtgccg 4860
 tcgccggagc tgtacacagt gatcctccgg ctggttgatg aggcggaacc ctattacgcg 4920
 tcggcgcggg tgggtctggc ggatctgcca ccgcgctgc cctggtccat cgccgccgcg 4980
 ctacggatct atcgcccat cgggtgcgc atccgcaaga gcgggcccga ggcctatcgc 5040
 cagcggatca gcacgtocaa ggctgccaag atcggcctgc tgggcgtcgg gggctgggat 5100
 gtcgcgcgat cacgcctgcc gggggcgggc gtgtcgcggc agggcctctg gaccggcccg 5160
 catcacgtct aggcgcgcgc ggcgtagggc agaaccctt ccagcagggc cgcgatttcc 5220

ggagcctgaa ggcgcttget ggcgagcatc gcgtccagtt gggcgcggtt ggcctcgtaa 5280
 tgacgggaca cgttctgcag gtctgacacg gccagaaggc cgcgcgcgcg gccggggggc 5340
 ggggcatcgc gaccggtatc cttgccaaagc gccgcctggt cgcgcacgac gtccagcagg 5400
 tcgtcatagg actggaacac ggggcccagc tgacggccaa agtcgatcat ctgggtctgc 5460
 tcctcggcgt cgaactcctt gatcacggcc agcatctcca gccggcgat gaacagcacg 5520
 ccggtcttca ggtcctgttc ctgttcgacc ccgcgcgctt tcttgccgcg gtgcagggtc 5580
 aggtcctggc cggcgacacg gccctgcggc ccaggggacc gcgacaggat ccgcaccagc 5640
 tgcgcccgca ccgtgcccga cgcgcgcgcg gcaccggcca gcaggggcat tgccctcggtg 5700
 atcaggggca tgccgcccag cacggcacgg ctttcgccat gcgccacatg ggtcgcgggc 5760
 cggccgcggc gcagcccggc atcgtccatg cagggcaggc cgtogaagat cagcgatgcg 5820
 gcatgcacca tctcgaccgc gcaggcgggc tcgacgatcg tgcgcgagc ccgcgccgag 5880
 gcctctgccg caagcagcat cagcatgccg cggaaccgcc tgcccgacga cagcgcgcca 5940
 tggctcatgg ccgcgcgag cggctgcgac acggcaccga atccctgggc gatctcctca 6000
 agtctggtct gcagaagggc ggcgtggatc ggggtgacgt ctgtctcat cagtgccttc 6060
 gcgcttgggt tctgacctgg cgggaaggc agggcggggc ggcaccccg gaccctcat 6120
 ccaccgtcaa cagtcccat gttggaacgg ttcacgccc attgcgagcc ttttcgacgg 6180
 cgacgcgggg tcgcgcggca atttgtcaa caaggtcagt gga 6223

<210> 5
 <211> 729
 <212> DNA
 <213> crtW gene

<400> 5
 atgagcgac atgccctgcc caaggcagat ctgaccgcca ccagcctgat cgtctcgggc 60
 ggcatcatcg ccgcgtggct ggcctgcat gtgcagcgc tgtggtttct ggacgcggcg 120
 gcgcatccca tcctggcgat cgcgaatttc ctggggctga cctggctgtc ggtcggctctg 180
 ttcttcatcg cgcagacgc gatgcacggg tcggtcgtgc cggggcgctc gcgcggcaat 240
 gcggcgatgg gccagctggt cctgtggctg tatgccgat tttcgtggcg caagatgatc 300
 gtcaagcaca tggcccatca ccgccatacc ggaaccgacg acgaccccg tttcgaccat 360
 ggcggcccg tccgctggta cgcgcgcttc atcggcacct atttcggctg gcgcgagggg 420
 ctgctgctgc ccgtcatcgt gacggtctat gcgctgatcc tgggggatcg ctggatgtac 480

gtggttcttct ggccgctgcc gtcgatcctg gcgtcgatcc agctgttcgt gttcggcacc 540
 tggtgcccgc accgccccgg ccacgacgcg ttcccggacc gccataatgc gcggtcgtcg 600
 cggatcagcg accccgtgtc gctgctgacc tgctttcact ttggtggtta tcatcacgaa 660
 caccacctgc acccgacggt gccttggtgg cgctgccca gcacccgcac caagggggac 720
 accgcatga 729

<210> 6
 <211> 242
 <212> PRT
 <213> crtW amino acid

<400> 6
 Met Ser Ala His Ala Leu Pro Lys Ala Asp Leu Thr Ala Thr Ser Leu
 1 5 10 15
 Ile Val Ser Gly Gly Ile Ile Ala Ala Trp Leu Ala Leu His Val His
 20 25 30
 Ala Leu Trp Phe Leu Asp Ala Ala Ala His Pro Ile Leu Ala Ile Ala
 35 40 45
 Asn Phe Leu Gly Leu Thr Trp Leu Ser Val Gly Leu Phe Phe Ile Ala
 50 55 60
 His Asp Ala Met His Gly Ser Val Val Pro Gly Arg Pro Arg Gly Asn
 65 70 75 80
 Ala Ala Met Gly Gln Leu Val Leu Trp Leu Tyr Ala Gly Phe Ser Trp
 85 90 95
 Arg Lys Met Ile Val Lys His Met Ala His His Arg His Thr Gly Thr
 100 105 110
 Asp Asp Asp Pro Asp Phe Asp His Gly Gly Pro Val Arg Trp Tyr Ala
 115 120 125
 Arg Phe Ile Gly Thr Tyr Phe Gly Trp Arg Glu Gly Leu Leu Leu Pro
 130 135 140
 Val Ile Val Thr Val Tyr Ala Leu Ile Leu Gly Asp Arg Trp Met Tyr
 145 150 155 160
 Val Val Phe Trp Pro Leu Pro Ser Ile Leu Ala Ser Ile Gln Leu Phe
 165 170 175
 Val Phe Gly Thr Trp Leu Pro His Arg Pro Gly His Asp Ala Phe Pro
 180 185 190
 Asp Arg His Asn Ala Arg Ser Ser Arg Ile Ser Asp Pro Val Ser Leu
 195 200 205

Leu Thr Cys Phe His Phe Gly Gly Tyr His His Glu His His Leu His
 210 215 220

Pro Thr Val Pro Trp Trp Arg Leu Pro Ser Thr Arg Thr Lys Gly Asp
 225 230 235 240

Thr Ala

<210> 7
 <211> 489
 <212> DNA
 <213> crtZ gene

<400> 7
 atgaccaatt tcctgatcgt cgtcgccacc gtgctggtga tggagttgac ggcctattcc 60
 gtccaccggtt ggatcatgca cggccccctg ggctggggct ggcacaagtc ccaccacgag 120
 gaacacgacc acgcgctgga aaagaacgac ctgtacggcc tggctctttgc ggtgatcgcc 180
 acggtgctgt tcacggtggg ctggatctgg gcgccggtcc tgtggtggat cgctttgggc 240
 atgaccgtct atgggctgat ctatttcgtc ctgcatgacg ggctggttca tcagcgctgg 300
 ccgttcgcgt atatcccgcg caagggctat gcccgccgcc tgtatcaggc ccaccgcctg 360
 caccacgcgg tcgagggacg cgaccattgc gtcagcttcg gcttcattcta tgcgcgcgcc 420
 gtcgacaagc tgaagcagga cctgaagacg tcgggcgtgc tgcgggccga ggcgcaggag 480
 cgcacgtga 489

<210> 8
 <211> 162
 <212> PRT
 <213> crtZ amino acid

<400> 8
 Met Thr Asn Phe Leu Ile Val Val Ala Thr Val Leu Val Met Glu Leu
 1 5 10 15
 Thr Ala Tyr Ser Val His Arg Trp Ile Met His Gly Pro Leu Gly Trp
 20 25 30
 Gly Trp His Lys Ser His His Glu Glu His Asp His Ala Leu Glu Lys
 35 40 45
 Asn Asp Leu Tyr Gly Leu Val Phe Ala Val Ile Ala Thr Val Leu Phe
 50 55 60
 Thr Val Gly Trp Ile Trp Ala Pro Val Leu Trp Trp Ile Ala Leu Gly
 65 70 75 80
 Met Thr Val Tyr Gly Leu Ile Tyr Phe Val Leu His Asp Gly Leu Val

85 90 95
 His Gln Arg Trp Pro Phe Arg Tyr Ile Pro Arg Lys Gly Tyr Ala Arg
 100 105 110
 Arg Leu Tyr Gln Ala His Arg Leu His His Ala Val Glu Gly Arg Asp
 115 120 125
 His Cys Val Ser Phe Gly Phe Ile Tyr Ala Pro Pro Val Asp Lys Leu
 130 135 140
 Lys Gln Asp Leu Lys Thr Ser Gly Val Leu Arg Ala Glu Ala Gln Glu
 145 150 155 160
 Arg Thr

<210> 9
 <211> 1161
 <212> DNA
 <213> crtY gene

<400> 9
 gtgacccatg acgtgctgct ggcaggggag ggccttgaga acgggctgat cgccctggcg 60
 ctgcgcgcgg cgcgccccga cctgcgggtg ctgctgctgg atcatgcggc gggaccgtca 120
 gacggccata cctggctcctg ccacgacccc gatctgtcgc cgcactggct ggcgcggtg 180
 aagcccctgc gccgcgccaa ctggcccgac caggaggtgc gctttcccg ccattgcccgg 240
 cggctggcca ccggttacgg gtcgctggac gggcgggcgc tggcggtatg ggtggcccg 300
 tcgggcgcgg agatccgctg gaacagcgac atcgccctgc tggatgaaca ggggcgacg 360
 ctgtcctgcg gcacccgat cgaggcgggc gcggtcctgg acgggcgcgg cgcgacggc 420
 tcgcggcatc tgaccgtggg ttccagaaa ttctggtggc tcgagatcga gaccgactgc 480
 cccacggcg tgccccgcc gatgatcatg gacgcgaccg tcaccagca ggacgggtac 540
 cgattcatct atctgctgcc cttctctccg acgcgcatcc tgatcgagga cactcgctat 600
 tccgatggcg gcaatctgga cgacgacgcg ctggcgggcg cgtccacga ctatgccgc 660
 cagcagggct ggaccggggc cgaggtccgg cggaacgcg gcatcctgcc cattgcgctg 720
 gccatgacg cggcgggctt ctgggcccga cacgcggagg ggctgttcc cgtgggactg 780
 cgcgcggggt tctttcaccc ggtcacggc tattcgctgc cctatgcggc gcaggtggcg 840
 gacgtggtgg cgggcctgtc cgggcccgc ggcaccgac cgctgcgcgg cgccatccgc 900
 gattacgcga tcgaccgggc acgccgtgac cgctttctgc gcctgctgaa ccggatgctg 960
 ttccgcggct gcgcgccccga ccggcgctat accctgctgc agcggttcta ccgcatgccg 1020

catggactga tcgaacggtt ctatgccggc cggctgagcg tggcggatca gctgcgcac 1080
 gtgaccggca agcctcccat tccccttggc acggccatcc gctgcctgcc cgaacgtccc 1140
 ctgctgaagg aaaacgcatg a 1161

<210> 10
 <211> 386
 <212> PRT
 <213> crtY amino acid

<400> 10
 Val Thr His Asp Val Leu Leu Ala Gly Ala Gly Leu Ala Asn Gly Leu
 1 5 10 15
 Ile Ala Leu Ala Leu Arg Ala Ala Arg Pro Asp Leu Arg Val Leu Leu
 20 25 30
 Leu Asp His Ala Ala Gly Pro Ser Asp Gly His Thr Trp Ser Cys His
 35 40 45
 Asp Pro Asp Leu Ser Pro His Trp Leu Ala Arg Leu Lys Pro Leu Arg
 50 55 60
 Arg Ala Asn Trp Pro Asp Gln Glu Val Arg Phe Pro Arg His Ala Arg
 65 70 75 80
 Arg Leu Ala Thr Gly Tyr Gly Ser Leu Asp Gly Ala Ala Leu Ala Asp
 85 90 95
 Ala Val Ala Arg Ser Gly Ala Glu Ile Arg Trp Asn Ser Asp Ile Ala
 100 105 110
 Leu Leu Asp Glu Gln Gly Ala Thr Leu Ser Cys Gly Thr Arg Ile Glu
 115 120 125
 Ala Gly Ala Val Leu Asp Gly Arg Gly Ala Gln Pro Ser Arg His Leu
 130 135 140
 Thr Val Gly Phe Gln Lys Phe Val Gly Val Glu Ile Glu Thr Asp Cys
 145 150 155 160
 Pro His Gly Val Pro Arg Pro Met Ile Met Asp Ala Thr Val Thr Gln
 165 170 175
 Gln Asp Gly Tyr Arg Phe Ile Tyr Leu Leu Pro Phe Ser Pro Thr Arg
 180 185 190
 Ile Leu Ile Glu Asp Thr Arg Tyr Ser Asp Gly Gly Asn Leu Asp Asp
 195 200 205
 Asp Ala Leu Ala Ala Ala Ser His Asp Tyr Ala Arg Gln Gln Gly Trp
 210 215 220
 Thr Gly Ala Glu Val Arg Arg Glu Arg Gly Ile Leu Pro Ile Ala Leu

225 230 235 240
 Ala His Asp Ala Ala Gly Phe Trp Ala Asp His Ala Glu Gly Pro Val
 245 250 255
 Pro Val Gly Leu Arg Ala Gly Phe Phe His Pro Val Thr Gly Tyr Ser
 260 265 270
 Leu Pro Tyr Ala Ala Gln Val Ala Asp Val Val Ala Gly Leu Ser Gly
 275 280 285
 Pro Pro Gly Thr Asp Ala Leu Arg Gly Ala Ile Arg Asp Tyr Ala Ile
 290 295 300
 Asp Arg Ala Arg Arg Asp Arg Phe Leu Arg Leu Leu Asn Arg Met Leu
 305 310 315 320
 Phe Arg Gly Cys Ala Pro Asp Arg Arg Tyr Thr Leu Leu Gln Arg Phe
 325 330 335
 Tyr Arg Met Pro His Gly Leu Ile Glu Arg Phe Tyr Ala Gly Arg Leu
 340 345 350
 Ser Val Ala Asp Gln Leu Arg Ile Val Thr Gly Lys Pro Pro Ile Pro
 355 360 365
 Leu Gly Thr Ala Ile Arg Cys Leu Pro Glu Arg Pro Leu Leu Lys Glu
 370 375 380
 Asn Ala
 385

<210> 11
 <211> 1506
 <212> DNA
 <213> crtI gene

<400> 11
 atgaacgccc attcgccgcg gcccaagacc gccatcgtga tcggcgagcag ctttggcggg 60
 ctggcccttg ccacccgcct gcagtcgcgc gccatcgcca ccaccctggt cgaggcccgg 120
 gacaagcccg gcgggcgcg cctatgtctg cagatcagc gccatgtctt cgacgcgggc 180
 ccgaccgtca tcaccgacc cgatgcgctc aaggagctgt gggcgctgac cgggcaggac 240
 atggcgcgcg acgtgacgt gatgccgtg tcgcccttct atcgactgat gtggccgggc 300
 gggaaggctt tcgattacgt gaacgaggcc gatcagctgg agcgccagat cgcccagttc 360
 aaccggagc acctggaagg ataccgcgc ttccgtgatt acggggagga ggtgtatcag 420
 gagggctacg tcaagctggg caccgtgcc ttctcaagc tggggcagat gctcaaggcc 480
 gcgcccgcgc tgatgaagct ggaggcctat aagtccgtcc atgccaaggt cgcgaccttc 540

atcaaggacc cctatctgcg gcaggcggtt tcgtatcaca cgctgctggt gggcggggaat 600
 cccttctcga ccagctcgat ctatgcgctg atccacgcgc tggagcgggc cggcggggtc 660
 tggttcgcca agggcggcac caaccagctg gtcgcgggca tggtcgcgct gttegaacgg 720
 cttggcgggc agatgatgct gaacgccaag gtcgcccga tcgagaccga gggcgcgcg 780
 accacgggcg tcaccctggc ggacggggcg tctttaagg cgcacatggt cgccagcaac 840
 ggcgacgtca tgcacaacta tcgcgacctg ctggggccaca cggcccgcgg gcagagccgc 900
 gcgaaatcgc tggaccgcaa gcgctggtcc atgtcggtgt tcgtgctgca tttcggtctg 960
 cgcgaggcgc ccaaggacat cgcgcacac accatcctgt tcggcccccg ctacaggag 1020
 ctggtcaacg agatcttcaa gggcccgaag ctggccgagg atttctcgct gtacctgcat 1080
 tcgccctgca cgaccgatcc ggacatggcg cctccgggca tgtccacgca ttacgtgctg 1140
 gccccgtgc cgcactctggg ccgcgccgag atogattggg cggtcgaggg gccgcgctat 1200
 gccgaccgca tcctggcgtc cctggaggag cggtgatcc cgaacctgcg cgccaacctg 1260
 accacgacgc gcattctcac gcccgccgat ttccgacgc aactgaacgc ccatcacggc 1320
 agcgccttct cggtcgagcc gacctgacg caatccgct ggttcgggc gcacaaccgc 1380
 gacaagacga tccgcaactt ctatctggtc ggcgcgggca cccatccggg cgcgggcatt 1440
 ccgggcgtcg tgggctcggc caaggccacg gccaggtga tgctgtccga cctggcgggc 1500
 gcatga 1506

<210> 12
 <211> 501
 <212> PRT
 <213> crtI amino acid

<400> 12
 Met Asn Ala His Ser Pro Ala Ala Lys Thr Ala Ile Val Ile Gly Ala
 1 5 10 15
 Gly Phe Gly Gly Leu Ala Leu Ala Ile Arg Leu Gln Ser Ala Gly Ile
 20 25 30
 Ala Thr Thr Leu Val Glu Ala Arg Asp Lys Pro Gly Gly Arg Ala Tyr
 35 40 45
 Val Trp His Asp Gln Gly His Val Phe Asp Ala Gly Pro Thr Val Ile
 50 55 60
 Thr Asp Pro Asp Ala Leu Lys Glu Leu Trp Ala Leu Thr Gly Gln Asp
 65 70 75 80
 Met Ala Arg Asp Val Thr Leu Met Pro Val Ser Pro Phe Tyr Arg Leu

85					90					95					
Met	Trp	Pro	Gly	Gly	Lys	Val	Phe	Asp	Tyr	Val	Asn	Glu	Ala	Asp	Gln
			100					105					110		
Leu	Glu	Arg	Gln	Ile	Ala	Gln	Phe	Asn	Pro	Asp	Asp	Leu	Glu	Gly	Tyr
		115					120					125			
Arg	Arg	Phe	Arg	Asp	Tyr	Ala	Glu	Glu	Val	Tyr	Gln	Glu	Gly	Tyr	Val
	130					135					140				
Lys	Leu	Gly	Thr	Val	Pro	Phe	Leu	Lys	Leu	Gly	Gln	Met	Leu	Lys	Ala
145					150					155					160
Ala	Pro	Ala	Leu	Met	Lys	Leu	Glu	Ala	Tyr	Lys	Ser	Val	His	Ala	Lys
				165					170					175	
Val	Ala	Thr	Phe	Ile	Lys	Asp	Pro	Tyr	Leu	Arg	Gln	Ala	Phe	Ser	Tyr
			180					185					190		
His	Thr	Leu	Leu	Val	Gly	Gly	Asn	Pro	Phe	Ser	Thr	Ser	Ser	Ile	Tyr
	195						200					205			
Ala	Leu	Ile	His	Ala	Leu	Glu	Arg	Arg	Gly	Gly	Val	Trp	Phe	Ala	Lys
	210					215					220				
Gly	Gly	Thr	Asn	Gln	Leu	Val	Ala	Gly	Met	Val	Ala	Leu	Phe	Glu	Arg
225				230						235					240
Leu	Gly	Gly	Gln	Met	Met	Leu	Asn	Ala	Lys	Val	Ala	Arg	Ile	Glu	Thr
				245					250					255	
Glu	Gly	Ala	Arg	Thr	Thr	Gly	Val	Thr	Leu	Ala	Asp	Gly	Arg	Ser	Leu
		260						265					270		
Arg	Ala	Asp	Met	Val	Ala	Ser	Asn	Gly	Asp	Val	Met	His	Asn	Tyr	Arg
		275					280					285			
Asp	Leu	Leu	Gly	His	Thr	Ala	Arg	Gly	Gln	Ser	Arg	Ala	Lys	Ser	Leu
	290					295					300				
Asp	Arg	Lys	Arg	Trp	Ser	Met	Ser	Leu	Phe	Val	Leu	His	Phe	Gly	Leu
305				310						315					320
Arg	Glu	Ala	Pro	Lys	Asp	Ile	Ala	His	His	Thr	Ile	Leu	Phe	Gly	Pro
			325					330					335		
Arg	Tyr	Arg	Glu	Leu	Val	Asn	Glu	Ile	Phe	Lys	Gly	Pro	Lys	Leu	Ala
			340					345					350		
Glu	Asp	Phe	Ser	Leu	Tyr	Leu	His	Ser	Pro	Cys	Thr	Thr	Asp	Pro	Asp
	355						360					365			
Met	Ala	Pro	Pro	Gly	Met	Ser	Thr	His	Tyr	Val	Leu	Ala	Pro	Val	Pro
	370					375					380				
His	Leu	Gly	Arg	Ala	Glu	Ile	Asp	Trp	Ala	Val	Glu	Gly	Pro	Arg	Tyr

385 390 395 400
 Ala Asp Arg Ile Leu Ala Ser Leu Glu Glu Arg Leu Ile Pro Asn Leu
 405 410 415
 Arg Ala Asn Leu Thr Thr Thr Arg Ile Phe Thr Pro Ala Asp Phe Ala
 420 425 430
 Ser Glu Leu Asn Ala His His Gly Ser Ala Phe Ser Val Glu Pro Ile
 435 440 445
 Leu Thr Gln Ser Ala Trp Phe Arg Pro His Asn Arg Asp Lys Thr Ile
 450 455 460
 Arg Asn Phe Tyr Leu Val Gly Ala Gly Thr His Pro Gly Ala Gly Ile
 465 470 475 480
 Pro Gly Val Val Gly Ser Ala Lys Ala Thr Ala Gln Val Met Leu Ser
 485 490 495
 Asp Leu Ala Gly Ala
 500

<210> 13
 <211> 915
 <212> DNA
 <213> crtB gene

<400> 13
 atgagcgatc tggctcctgac ctcgaccgag gcgatcacc aagggtcgca aagctttgcc 60
 acggcggcca agctgatgcc gccgggcac cgcgacgaca cggatgatgct ctatgcctgg 120
 tgccgccacg cggatgacgt gatcgacggt caggccctgg gcagccgccc cgaggcgggtg 180
 aacgacccgc aggcgcggct ggacggcctg cgcgtcgaca cgctggcggc cctgcagggc 240
 gacgggtccg tgaccccgcc ctttgccgcg ctgcgcgcgg tggcgcggcg gcatgatattc 300
 ccgcaggcct ggcccatgga cctgatcgaa ggcttcgcga tggatgtcga ggcgcgcgac 360
 tatcgacgc tggatgacgt gctggaatat tcctatcacg tcgcaggcat cgtcggcgtg 420
 atgatggccc gcgtgatggg cgtgcgcgac gatcctgtcc tggaccgcgc ctgcgacctg 480
 gggctggcgt tccagctgac caacatcgcg cgcgacgtga tcgacgatgc gcgcacggg 540
 cgggtgctatc tgccggggga ctggctggac caggcgggcg cgcggatcga cgggccgggtg 600
 ccgtcgccgg agctgtacac agtgatcctc cggctgttgg atgaggcgga accctattac 660
 gcgtcggcgc ggggtgggtct ggcggatctg ccaccgcgct gcgcctggtc catcgccgcc 720
 gcgctacgga tctatcgcg catcgggctg cgcacccgca agagcgggccc gcaggcctat 780
 cgccagcgga tcagcacgtc caaggctgcc aagatcggcc tgctgggcgt cgggggctgg 840

gatgtcgcgc gatcacgcct gccgggggcg ggcgtgtcgc ggcagggcct ctggaccggg 900
ccgcatcacg tctag 915

<210> 14
<211> 304
<212> PRT
<213> crtB amino acid

<400> 14
Met Ser Asp Leu Val Leu Thr Ser Thr Glu Ala Ile Thr Gln Gly Ser
1 5 10 15
Gln Ser Phe Ala Thr Ala Ala Lys Leu Met Pro Pro Gly Ile Arg Asp
20 25 30
Asp Thr Val Met Leu Tyr Ala Trp Cys Arg His Ala Asp Asp Val Ile
35 40 45
Asp Gly Gln Ala Leu Gly Ser Arg Pro Glu Ala Val Asn Asp Pro Gln
50 55 60
Ala Arg Leu Asp Gly Leu Arg Val Asp Thr Leu Ala Ala Leu Gln Gly
65 70 75 80
Asp Gly Pro Val Thr Pro Pro Phe Ala Ala Leu Arg Ala Val Ala Arg
85 90 95
Arg His Asp Phe Pro Gln Ala Trp Pro Met Asp Leu Ile Glu Gly Phe
100 105 110
Ala Met Asp Val Glu Ala Arg Asp Tyr Arg Thr Leu Asp Asp Val Leu
115 120 125
Glu Tyr Ser Tyr His Val Ala Gly Ile Val Gly Val Met Met Ala Arg
130 135 140
Val Met Gly Val Arg Asp Asp Pro Val Leu Asp Arg Ala Cys Asp Leu
145 150 155 160
Gly Leu Ala Phe Gln Leu Thr Asn Ile Ala Arg Asp Val Ile Asp Asp
165 170 175
Ala Arg Ile Gly Arg Cys Tyr Leu Pro Gly Asp Trp Leu Asp Gln Ala
180 185 190
Gly Ala Arg Ile Asp Gly Pro Val Pro Ser Pro Glu Leu Tyr Thr Val
195 200 205
Ile Leu Arg Leu Leu Asp Glu Ala Glu Pro Tyr Tyr Ala Ser Ala Arg
210 215 220
Val Gly Leu Ala Asp Leu Pro Pro Arg Cys Ala Trp Ser Ile Ala Ala
225 230 235 240

Ala Leu Arg Ile Tyr Arg Ala Ile Gly Leu Arg Ile Arg Lys Ser Gly
 245 250 255

Pro Gln Ala Tyr Arg Gln Arg Ile Ser Thr Ser Lys Ala Ala Lys Ile
 260 265 270

Gly Leu Leu Gly Val Gly Gly Trp Asp Val Ala Arg Ser Arg Leu Pro
 275 280 285

Gly Ala Gly Val Ser Arg Gln Gly Leu Trp Thr Arg Pro His His Val
 290 295 300

<210> 15
 <211> 882
 <212> DNA
 <213> crtE gene

<400> 15
 atgagacgag acgtcaaccc gatccacgcc acccttctgc agaccagact tgaggagatc 60
 gccagggat tcggtgccgt gtcgcagccg ctggcgcgcg ccatgagcca tggcgcgctg 120
 tcgtcgggca ggcggttccg cggcatgctg atgotgcttg cggcagaggc ctcgggcggg 180
 gtctgcgaca cgatcgtcga cgccgcctgc gcggtcgaga tgggtcatgc cgcacgctg 240
 atcttcgacg acctgccctg catggacgat gccgggctgc gccgcggccg gcccgcgacc 300
 catgtggcgc atggcgaaag ccgtgccgtg ctggggcgga tcgccctgat caccgaggca 360
 atggccctgc tggccggtgc gcgcggcgcg tcgggcacgg tgcgggcgca gctggtgcgg 420
 atcctgtcgc ggtccctggg gccgcagggc ctgtgcgccg gccaggacct ggacctgcac 480
 gcggccaaga acggcgcggg ggtcgaacag gaacaggacc tgaagaccgg cgtgctgttc 540
 atcgccgggc tggagatgct ggccgtgac aaggagttcg acgccgagga gcagaccag 600
 atgatcgact ttggccgtca gctgggcccgc gtgttccagt cctatgacga cctgctggac 660
 gtcgtgggcg accaggcggc gcttggcaag gataccggtc gcgatgccgc ggcccccggc 720
 ccgcggcgcg gccttctggc cgtgtcagac ctgcagaacg tgtcccgtca ttacgaggcc 780
 agccgcgccc aactggacgc gatgctgcgc agcaagcgcc ttcaggctcc ggaaatcgcg 840
 gccctgctgg aacgggttct gccctacgcc gcgcgcgcct ag 882

<210> 16
 <211> 293
 <212> PRT
 <213> crtE amino acid

```

<400>    16
Met Arg Arg Asp Val Asn Pro Ile His Ala Thr Leu Leu Gln Thr Arg
 1          5          10          15

Leu Glu Glu Ile Ala Gln Gly Phe Gly Ala Val Ser Gln Pro Leu Gly
      20          25          30

Ala Ala Met Ser His Gly Ala Leu Ser Ser Gly Arg Arg Phe Arg Gly
      35          40          45

Met Leu Met Leu Leu Ala Ala Glu Ala Ser Gly Gly Val Cys Asp Thr
      50          55          60

Ile Val Asp Ala Ala Cys Ala Val Glu Met Val His Ala Ala Ser Leu
      65          70          75          80

Ile Phe Asp Asp Leu Pro Cys Met Asp Asp Ala Gly Leu Arg Arg Gly
      85          90          95

Arg Pro Ala Thr His Val Ala His Gly Glu Ser Arg Ala Val Leu Gly
      100          105          110

Gly Ile Ala Leu Ile Thr Glu Ala Met Ala Leu Leu Ala Gly Ala Arg
      115          120          125

Gly Ala Ser Gly Thr Val Arg Ala Gln Leu Val Arg Ile Leu Ser Arg
      130          135          140

Ser Leu Gly Pro Gln Gly Leu Cys Ala Gly Gln Asp Leu Asp Leu His
      145          150          155          160

Ala Ala Lys Asn Gly Ala Gly Val Glu Gln Glu Gln Asp Leu Lys Thr
      165          170          175

Gly Val Leu Phe Ile Ala Gly Leu Glu Met Leu Ala Val Ile Lys Glu
      180          185          190

Phe Asp Ala Glu Glu Gln Thr Gln Met Ile Asp Phe Gly Arg Gln Leu
      195          200          205

Gly Arg Val Phe Gln Ser Tyr Asp Asp Leu Leu Asp Val Val Gly Asp
      210          215          220

Gln Ala Ala Leu Gly Lys Asp Thr Gly Arg Asp Ala Ala Ala Pro Gly
      225          230          235          240

Pro Arg Arg Gly Leu Leu Ala Val Ser Asp Leu Gln Asn Val Ser Arg
      245          250          255

His Tyr Glu Ala Ser Arg Ala Gln Leu Asp Ala Met Leu Arg Ser Lys
      260          265          270

Arg Leu Gln Ala Pro Glu Ile Ala Ala Leu Leu Glu Arg Val Leu Pro
      275          280          285

Tyr Ala Ala Arg Ala
      290

```

<210> 17
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> forward primer for crt gene

<400> 17
gttccacgac tggggcatc 19

<210> 18
<211> 28
<212> DNA
<213> Artificial Sequence

<220>
<223> reverse primer for crt gene

<400> 18
tccactgacc ttgttgaca aattgccg 28

INTERNATIONAL SEARCH REPORT

International application No.

PCT/KR2004/000752

A. CLASSIFICATION OF SUBJECT MATTER**IPC7 C12N 1/20**

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
Korean patents and applications for inventions since 1975Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
PubMed, NCBI, KIPASS, Delphion, CA, PAJ**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	NCBI accession No. D58420 (28 August 2002)	1,2,3
X	NCBI accession No. Y15112 (15 September 1999)	1,4-8, 10-17
X	NCBI accession No. BAA09591 (28 August 2002)	9
X	NCBI accession No. CAB56060 (15 September 1999)	9
X	NCBI accession No. BAA09593 (28 August 2002)	9
X	NCBI accession No. CAB56062 (15 September 1999)	9
X	NCBI accession No. CAB56063 (15 September 1999)	9
X	NCBI accession No. CAB56064 (15 September 1999)	9



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

28 JULY 2004 (28.07.2004)

Date of mailing of the international search report

29 JULY 2004 (29.07.2004)

Name and mailing address of the ISA/KR

Korean Intellectual Property Office
920 Dunsan-dong, Seo-gu, Daejeon 302-701,
Republic of Korea

Facsimile No. 82-42-472-7140

Authorized officer

KIM, Jae Hyun

Telephone No. 82-42-481-5591



INTERNATIONAL SEARCH REPORT

International application No.

PCT/KR2004/000752

Box No. I Nucleotide and/or amino acid sequence(s) (Continuation of item 1.b of the first sheet)

1. With regard to any nucleotide and/or amino acid sequence disclosed in the international application and necessary to the claimed invention, the international search was carried out on the basis of:

a. type of material



a sequence listing



table(s) related to the sequence listing

b. format of material



in written format



in computer readable form

c. time of filing/furnishing



contained in the international application as filed



filed together with the international application in computer readable form



furnished subsequently to this Authority for the purposes of search

2. ☐ In addition, in the case that more than one version or copy of a sequence listing and/or table relating thereto has been filed or furnished, the required statements that the information in the subsequent or additional copies is identical to that in the application as filed or does not go beyond the application as filed, as appropriate, were furnished.

3. Additional comments: